

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

The topology and drivers of ant-symbiont networks across Europe

Parmentier, Thomas; DE LAENDER, Frederik; Bonte, Dries

Published in:
Biological Reviews

DOI:
[10.1111/brv.12634](https://doi.org/10.1111/brv.12634)

Publication date:
2020

Document Version
Peer reviewed version

[Link to publication](#)

Citation for published version (HARVARD):

Parmentier, T, DE LAENDER, F & Bonte, D 2020, 'The topology and drivers of ant-symbiont networks across Europe', *Biological Reviews*, vol. 95, no. 6, pp. 1664-1688. <https://doi.org/10.1111/brv.12634>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

The topology and drivers of ant–symbiont networks across Europe

Thomas Parmentier^{1,2,*}, Frederik de Laender^{2,†} and Dries Bonte^{1,†}

¹*Terrestrial Ecology Unit (TEREC), Department of Biology, Ghent University, K.L. Ledeganckstraat 35, B-9000 Ghent, Belgium*

²*Research Unit of Environmental and Evolutionary Biology, Namur Institute of Complex Systems, and Institute of Life, Earth, and the Environment, University of Namur, Rue de Bruxelles 61, 5000 Namur, Belgium*

* Author for correspondence at address 1 (E-mail: Thomas.Parmentier@ugent.be; Tel.: +32 (0)9 264 52 60).

† These authors contributed equally.

ABSTRACT

Intimate associations between different species drive community composition across ecosystems. Understanding the ecological and evolutionary drivers of these symbiotic associations is challenging because their structure eventually determines stability and resilience of the entire species network. Here, we compiled a detailed database on naturally occurring ant–symbiont networks in Europe to identify factors that affect symbiont network topology. These networks host an unrivalled diversity of macrosymbiotic associations, spanning the entire mutualism–antagonism continuum, including: (1) myrmecophiles –

commensalistic and parasitic arthropods; (2) trophobionts – mutualistic aphids, scale insects, planthoppers and caterpillars; (3) social parasites – parasitic ant species; (4) parasitic helminths; and (5) parasitic fungi. We dissected network topology to investigate what determines host specificity, symbiont species richness, and the capacity of different symbiont types to switch hosts.

We found 722 macrosymbionts (multicellular symbionts) associated with European ants. Symbiont type explained host specificity and the average relatedness of the host species. Social parasites were associated with few hosts that were phylogenetically highly related, whereas the other symbiont types interacted with a larger number of hosts across a wider taxonomic distribution. The hosts of trophobionts were the least phylogenetically related across all symbiont types. Colony size, host range and habitat type predicted total symbiont richness: ant hosts with larger colony size, a larger distribution range or with a wider habitat range contained more symbiont species. However, we found that different sets of host factors affected diversity in the different types of symbionts. Ecological factors, such as colony size, host range and niche width predominantly determined myrmecophile species richness, whereas host phylogeny was the most important predictor of mutualistic trophobiont, social parasite and parasitic helminth species richness. Lastly, we found that hosts with a common biogeographic history support a more similar community of symbionts. Phylogenetically related hosts also shared more trophobionts, social parasites and helminths, but not myrmecophiles.

Taken together, these results suggest that ecological and evolutionary processes structure host specificity and symbiont richness in large-scale ant–symbiont networks, but these drivers may shift in importance depending on the type of symbiosis. Our findings highlight the potential of well-characterized bipartite networks composed of different types of symbioses to identify candidate processes driving community composition.

50 *Key words:* ant guests, commensalism, ecological network, Formicidae, host–parasite
51 network, host switching, inquiline, microcosm, mutualism, parasite species richness.
52

53 CONTENTS

54	I. Introduction.....	3
55	II. Materials and methods	8
56	(1) Ant symbionts.....	8
57	(2) Ant–symbiont data set compilation	9
58	(3) Host specificity in different symbiont types.....	10
59	(4) Predictors of symbiont diversity in European ants	12
60	(5) Predictors of symbiont sharing in European ants	15
61	III. Results.....	16
62	(1) Ant–symbiont networks display a diversity of species interactions.....	16
63	(2) Host specificity in different symbiont types.....	18
64	(3) Predictors of symbiont diversity in European ants	18
65	(4) Predictors of symbiont sharing in European ants	20
66	IV. Discussion	21
67	(1) Characterization of the European ant–symbiont network	21
68	(2) Host specificity in different symbiont types.....	22
69	(3) Predictors of symbiont diversity in European ants	25
70	(4) Predictors of symbiont sharing in European ants	27
71	V. Outstanding questions	29
72	VI. Conclusions.....	31
73	VII. Acknowledgements.....	32
74	VIII. References	32

75	IX. Supporting information	73
----	----------------------------------	----

76

77 **I. INTRODUCTION**

78 Close associations between different species, known as symbioses, are crucial components of
79 communities in all ecosystems. These intricate associations display a high diversity, ranging
80 from mutually beneficial partnerships to parasitic interactions in which one species exploits
81 another (Paracer & Ahmadjian, 2000). Interactions among species are central to ecological
82 and evolutionary dynamics in assemblages of species that belong to different guilds and
83 trophic levels. They are thus essential elements of the ‘entangled bank’ – Darwin’s metaphor
84 for the complexity and connectedness of natural systems (Darwin, 1859) – and can give rise
85 to important stabilising feedbacks that eventually maintain diversity and ecosystem
86 functioning (Thrall *et al.*, 2007; Bastolla *et al.*, 2009). To date, insights are derived from
87 theory and empiricism focusing on either competition and predator–prey interactions (e.g.
88 Hairston, Smith & Slobodkin, 1960; Pimm, 1979; Tilman, 1982) or more recently mutualistic
89 interactions (Bascompte, Jordano & Olesen, 2006; Bascompte & Jordano, 2007; Barabás,
90 D’Andrea & Stump, 2018; Johnson & Bronstein, 2019; Spaak & De Laender, 2020).
91 Communities contain a wide variety of interactions, rendering the ecological network
92 extremely complex (Newman, 2003). One important feature of complex systems is the
93 presence of properties that emerge from interactions among the specific components of the
94 system (Solé & Bascompte, 2006). Typically, these emergent properties result from the
95 interaction between different network components across time and space and are difficult to
96 predict from the specific (isolated) pairwise interactions (Vázquez, Chacoff & Cagnolo, 2009;
97 Grilli *et al.*, 2017). Generally speaking, modular networks that are characterized by a high
98 connectance tend to be more robust to species loss, and are less affected by disturbance (Solé
99 & Bascompte, 2006; Olesen *et al.*, 2007). While theoretical progress has been made (Solé &

100 Montoya, 2001), the field is suffering from a lack of comprehensive and manageable
101 empirical systems. This limits empirical tests of theory (Valdovinos, 2019) and thus hampers
102 the development of effective management tools to safeguard the biodiversity and ecosystem
103 functions of natural ecosystems (De Laender *et al.*, 2016).

104 The study and interpretation of the drivers and consequences of the topology in host–
105 symbiont networks sheds light on the evolution of the strategies and traits of hosts and
106 symbionts. Moreover, this network approach may unravel general rules in community
107 assembly processes which can be compared across different types of symbionts and different
108 systems. While network studies at local scales will be informative on the local community
109 assembly processes, those reconstructed at regional or global scales will allow inference of
110 macro-ecological and evolutionary processes (Trøjelsgaard & Olesen, 2013).

111 In contrast to trophic networks, bipartite host–symbiont networks contain different kinds of
112 links, with interactions between hosts and symbionts ranging from antagonistic to mutualistic
113 (Ings *et al.*, 2009). Examples of such networks include plant–mycorrhiza and host–
114 microbiome associations. The complexity of these networks is enormous, and their
115 description is merely based on one interaction type, either antagonistic or mutualistic,
116 although theory predicts that the diversity of interaction types may be essential for community
117 stability (Fontaine *et al.*, 2011; Mougi & Kondoh, 2012; Allesina & Tang, 2015). The
118 topology of bipartite host–symbiont networks can be dissected by adopting two different
119 perspectives, i.e. that of each of the individual sets of species (hosts and symbionts). Asking
120 what factors cause a given topology is equivalent to asking, for each of the sets, what explains
121 the number of links per species and the specificity of these links, i.e. how the links are
122 distributed among species from the focal set. An example of this approach is found in studies
123 on predator–prey networks, where average web vulnerability (i.e. the average number of

predators per prey) and generality (i.e. the average number of prey eaten per predator) link the specificity of the two interacting species sets (Schoener, 1989).

A determining feature of the ecology of symbionts is host specificity, which can be quantified in host–symbiont networks by the number of links departing from a symbiont node. Yet, a measure of host specificity should ideally consider the relatedness of the targeted host species as well (Poulin & Mouillot, 2003). Generalist symbionts target multiple, unrelated host species and may gain low to high benefits in any host. Specialist symbionts, in turn, engage with one or only a few related species, and achieve high benefits with their hosts by resorting to advanced morphological, physiological and behavioural adaptations (Bronstein, Alarcón & Geber, 2006; Thrall *et al.*, 2007). Their strong specialization, however, is offset by lower population densities and higher extinction risks due to the lower availability of hosts. Several studies in host–symbiont systems clearly found that host specificity is tightly linked with fundamental ecological processes and evolutionary history. Typically, host specificity is different among cohabiting symbionts, as for example demonstrated in parasites of primates (Pedersen *et al.*, 2005) and in parasitic mites on mussels (Edwards & Malcolm, 2006). A study on moths and plants indicated that host specificity can be dependent on the type of symbiotic interaction, with pollinating moths being more specific than their parasitic leaf-feeding relatives (Kawakita *et al.*, 2010).

From the perspective of the host, it is fundamental to understand the ecological, evolutionary and environmental drivers that promote the number of associated symbionts, i.e. the number of links departing from a host node to symbiont nodes. Studies on different host–symbiont systems reported multiple host variables which correlate with parasite species richness. Generally, the makeup of symbiont communities is orchestrated by both ecological and evolutionary host factors. Body size has been identified as a key ecological factor that favours species richness (Lindenfors *et al.*, 2007) and species interactions (Werner & Gilliam, 1984).

Host species may be conceptualized as island habitats for symbionts and, in line with island theory (MacArthur & Wilson, 1967), larger host species tend to support more symbionts (Kuris, Blaustein & Alió, 1980). Symbiont richness is also expected to increase with other ecological factors such as the number and variety of microhabitats offered by the host, host longevity, host range, and interaction probability with other host species (Kamiya *et al.*, 2014; Stephens *et al.*, 2016). Evolutionary processes may affect symbiont species richness in different ways. Related hosts often show traits that are correlated throughout evolution (phylogenetic correlation) which lead to similar values in species richness. However, related host species may have diverged with time, whether or not in a common spatial evolutionary ancestry (biogeography), but may still attract a similar fauna of symbionts as unrelated host species with a similar ecology (Poulin, 1995).

Another pattern that emerges in host–symbiont networks is the sharing/transmission of symbiont species across host species. The degree of symbiont sharing is vital as symbiont transmission can connect eco-evolutionary dynamics across hosts as a result of rapid symbiont spread in host populations [e.g. Jaenike *et al.*, 2010; Himler *et al.* (2011) in endosymbionts]. While little is known about the proximate mechanisms by which single symbionts switch between hosts, we can anticipate that host species with similar ecological niches and/or a shared evolutionary history tend to have similar symbiont communities. The pervasive effect of phylogenetic relatedness on symbiont sharing has for example been demonstrated in bat parasites (Luis *et al.*, 2015) and in plant–mycorrhiza (Veresoglou & Rillig, 2014).

Ant–symbiont networks are ideally suited to study which factors drive bipartite network topology (Ivens *et al.*, 2016). The diversity of symbiotic associations found in ants (Kistner, 1982; Hölldobler & Wilson, 1990; Rettenmeyer *et al.*, 2010; Parmentier, 2020) is thought to be promoted by their omnipresence in terrestrial ecosystems, their stable and climate-

controlled nest fortresses and the high number of available resources in the nest (Kronauer & Pierce, 2011). Ants interact with different types of symbionts spanning the entire parasitism–mutualism gradient. They include parasitic ants, different groups of arthropods living in the nests, mutualistic aphids and caterpillars, endoparasitic helminths, plants, bacteria and fungi (Hölldobler & Wilson, 1990). Therefore, they are promising systems to study different interaction types within a single biological system (Fontaine *et al.*, 2011).

Ant–symbiont networks that have been studied recently typically deal with local interaction networks and mostly focus on one kind of symbiotic interaction in isolation, such as mutualistic plant–ant networks (Guimarães *et al.*, 2006; Blüthgen *et al.*, 2007; Dáttilo, Guimarães & Izzo, 2013; Cagnolo & Tavella, 2015), mutualistic aphid–ant networks (Ivens *et al.*, 2018) or parasite–ant networks (Elizalde *et al.*, 2018). Some studies have already covered different types of symbiotic interactions in a local network (Pérez-Lachaud & Lachaud, 2014; Rocha, Lachaud & Pérez-Lachaud, 2020) and a recent study tested different types of ant–symbiont interactions on a large scale (Glasier, Poore, & Eldridge, 2018). However, the latter study only included a limited set of interaction types and pooled interactions of well-studied bioregions with those of very poorly studied regions.

Here, we ask what factors explain the topology of ant–symbiont networks across Europe. We firstly provide a quantitative and systematic meta-analysis of the diversity of European ant–symbiont interactions. By adopting the symbiont perspective, we test the hypothesis that the type of symbiosis explains the number and identity of their host species (host specificity). More specifically, we expect that parasitic ants are more specific than the other types of symbionts. Secondly, we follow a trait-based host perspective to identify the major drivers that promote the diversity of ant–symbiont interactions and facilitate symbiont sharing. We test the hypothesis that the number of symbionts with which an ant species interacts and the number of symbionts it shares with other ant hosts depend on ecological factors (colony size,

nest type, distribution, habitat, degree of sympatry, worker size) and evolutionary drivers (phylogeny, biogeography) associated with the host species.

II. MATERIALS AND METHODS

(1) Ant symbionts

Symbionts are species that engage in a close association with a host species on which they may have beneficial, neutral or adverse effects. We limited our analyses to Europe (excluding the Canary Islands and Madeira), as knowledge of ant–symbiont interactions on other continents is extremely fragmentary and poorly understood. Moreover myriad unknown symbionts presumably await discovery and description in these continents (Parmentier, 2020). By contrast, a firm body of knowledge on the distribution and diversity of ant symbionts in Europe has been recorded and has grown steadily from a long tradition of studying ant symbionts since the end of the 19th century (Wasmann, 1894; Janet, 1897). We focused on all types of macrosymbionts (multicellular organisms) associated with European ants. We did not include microsymbionts (unicellular microorganisms such as bacteria), as only a few case studies are available. In addition, these symbionts differ completely in their strategies and dynamics. Depending on the intimacy of the relationship between ants and symbionts, we can distinguish obligate and facultative interactions. An obligate interaction occurs when a symbiont lives permanently inside or near an ant nest. Obligate symbionts depend completely on ants and cannot be found without them. Facultative myrmecophiles may associate with ants, but regularly (or mostly) occur without ants. This study focuses on obligate symbionts. We categorized five types of symbionts: (1) myrmecophiles; (2) trophobionts; (3) social parasites; (4) helminths; and (5) fungi (Table 1). Myrmecophiles were further subdivided into three functional groups: unspecialized myrmecophiles, specialized myrmecophiles and myrmecophilous parasitoids (Table 1).

Plants engaging in mutualistic relationships (e.g. myrmecochory) were not included in our analyses. Contrary to the tropics, ant–plant relationships tend to be loose in Europe and are at best facultative (Rico-Gray & Oliveira, 2007).

(2) Ant–symbiont data set compilation

We compiled documented ant–symbiont interactions in Europe. Our database of ant–symbiont interactions was assembled from 269 published references, including faunistic notes, research articles, reviews and books. In a first round of searches, we scanned reference works (e.g. Wasmann, 1894; Donisthorpe, 1927; Evans & Till, 1966; Uppstrom, 2010; Tykarski, 2017; Molero-Baltanás *et al.* 2017) for associations between ant hosts and symbionts. Next, we searched for ant–symbiont interactions *via Google Scholar* using the terms: “myrmecophile” or “ant associate” or “inquiline” or “ant guest” or “ant symbiont”. We also found host–symbiont interactions within the reference lists of the retrieved publications. In a second phase, each symbiont occurring in Europe was searched by its Latin binomial name and its common taxonomic synonyms combined with a search string with the names of all ant genera ($N = 56$; AntWiki, 2019) found in Europe (for example “*Phengaris alcon*” AND *Acropyga* OR *Anochetus* OR *Aphaenogaster* OR *Bothriomyrmex* OR *Camponotus* OR ...) using *Google Scholar*. We chose *Google Scholar* over ISI Web of Science, as the latter does not retrieve faunistic notes or other types of grey literature. We omitted symbionts from our data set when they were reported not to be associated with ants. Note that some species with a poorly studied biology, such as parasitoid wasps and mites, were included in our data set, although they may not be completely dependent on ants. Ultimately, we obtained a binary host–symbiont matrix (see online Supporting information Appendix S1, references used to compile this table are listed in Appendix S2) filled with interactions (1) and non-interactions (0) between ants (columns) and symbionts (rows). We included some references on ant–

trophobiont interactions reported in the non-European part of Russia (e.g. Novgorodova, 2005) to increase the relatively modest number of reported interactions in this type of association. The reported ants and trophobionts in these references have a widespread Palearctic distribution and they are expected to interact in Europe as well.

(3) Host specificity in different symbiont types

We first determined the host range of the different symbiont types. A second analysis compared host relatedness across symbiont types. We used two approaches to estimate host relatedness: taxonomic and phylogenetic relatedness.

In the first analysis, we compared the number of host species across seven different types of host symbionts, i.e. unspecialized myrmecophiles, specialized myrmecophiles, myrmecophilous parasitoids, trophobionts, social parasites, parasitic fungi and helminths (Table 1). Symbionts with hosts only identified at the genus level were not included in all subsequent analyses. Note that we found evidence of one mutualistic fungus (*Cladosporium myrmecophilum*) (Table 1), but we only analysed parasitic fungi to ensure a coherent functional group of symbionts. Studies are uneven across symbionts, potentially meaning that better studied symbionts have a higher number of recorded host species. To account for differences in sampling effort, we therefore first performed a regression of the total number of host species against the $(\ln+1)$ -transformed number of *Google Scholar* hits for the binomial species name (and commonly used synonyms) of the symbionts. The residuals of this regression were not normally distributed. Therefore we used a non-parametric Kruskal-Wallis test to compare these residuals across symbiont types. This test was followed by *post-hoc* Dunn tests with the Benjamini–Hochberg adjustment for multiple testing.

In the second analysis, we compared the relatedness of targeted host species across the seven different symbiont types. For each symbiont, we estimated the average taxonomic distance

between the different hosts by using the specificity index, S_{TD} proposed by Poulin & Mouillot (2003). Host ant species (all ants belong to the family Formicidae) were classified following Linnaean classification into subfamilies, tribes, genera and species groups/subgenera. The taxonomic distance between two hosts is then defined as the number of hierarchical steps that are needed to reach a common node in the taxonomic tree. The taxonomic distance between two species of the same subgenus/species group equals 1; the distance between two species of the same genus, but from a different subgenus/species group equals 2. A distance of 3 or 4 was assigned to a pair of species belonging to the same tribe or subfamily, respectively. A taxonomic distance of 5 is reached between two ant host species from different subfamilies (largest possible distance). S_{TD} was estimated by averaging the taxonomic distance across all pairs of host species. However, S_{TD} cannot be calculated for symbionts with a single host species, although this can be addressed by excluding these ‘singletons’ from the analyses (Poulin & Mouillot, 2005). The single host species for many of our symbionts (e.g. mites) likely reflects undersampling, rather than true specificity to a single host species, because even extremely specialized species (e.g. *Microdon*, *Claviger*, social parasites) often have more than one host species. Since these singletons were equally distributed over all groups of symbionts, we argue they could be omitted from further analyses; a total of 392 symbionts were retained in our analysis. Note that the S_{TD} is not affected by uneven sampling when dealt with in this way. We modelled the S_{TD} as a response variable against the predictor symbiont type with a non-parametric Kruskal-Wallis test, followed by *post-hoc* Dunn tests with the Benjamini–Hochberg adjustment for multiple testing. Alternatively, we can assign a default taxonomic distance of 1 to the singletons, which is the lowest possible value (Poulin & Mouillot, 2005). We repeated the analysis with this approach and compared it with the analysis without singletons.

Next, we compared the relatedness of the targeted host species across the symbiont types with a phylogenetic- instead of a taxonomic-distance matrix. The phylogenetic-distance matrix was based on the phylogenetic tree of European ants (Arnan, Cerdá & Retana, 2017). Distances between species were estimated by node count (number of nodes along the branches of the tree from one species to another) and were retrieved using Mesquite v.3.5. Phylogenetic distances are more accurate than taxonomic distances to assess relatedness, but unfortunately we do not possess phylogenetic information at the species level for all ants in our data set (the phylogeny was known for 108 out of 181 ant species). We decided to exclude the 73 ant species without phylogenetic information and their interactions with their symbionts from subsequent analyses. We believe that this is acceptable as the retained 108 species cover 87.8% of the interactions in our host species–symbiont data set. In addition, symbionts that interacted with only one ant species were omitted, as no specificity index could be calculated for these species. Ultimately, we retained 362 symbiont species and 108 ant species in this analysis (host–symbiont matrix in Appendix S3). As the residuals meet the assumptions of a general linear model (GLM), we ran a GLM using phylogenetic distance as the response variable and symbiont type as predictor. Significance of the model was assessed with a likelihood ratio test (ANOVA function implemented in the package car), pairwise *post-hoc* tests were conducted with the Benjamini–Hochberg adjustment for multiple testing.

(4) Predictors of symbiont diversity in European ants

A central question in this study is why some ant species host more symbionts than other ant species. Therefore, we first compiled for the European ant species several predictors based on Arnan *et al.* (2017), Boulay *et al.* (2017), Seifert (2007) and AntWiki (2019). We selected traits of the host that could affect symbiont diversity. These predictors were colony size (number of workers), average worker size (mm), nest type [levels: (a) arboricolous; (b)

323 diverse; (c) soil; and (d) organic mound] and phylogeny as a proxy for trait similarity, and
 324 factors reflecting differences in the functional role of the host species, including habitat
 325 [levels: (a) eurytope: – a broad range of niches, including anthropogenic habitats; (b) open;
 326 (c) open/sylvicolous; and (d) sylvicolous], distribution range, the number of sympatric ants
 327 and biogeographic region [levels: (a) atlantic; (b) boreo-mountain; (c) continental; (d)
 328 Mediterranean; and (e) wide-ranging]. Ants were assigned to the biogeographical region
 329 where they were found proportionally the most in sampled biogeographical regions based on
 330 the observations of Arnan *et al.* (2017). If the proportional occurrence in the most preferred
 331 region was less than double the proportional occurrence in another region, the ant species was
 332 grouped under the ‘wide-ranging’ category. We also estimated the distribution range [the
 333 number of countries where the host species has been reported, based on records on AntWiki
 334 (2019)], the number of sympatric ants [number of ant species with symbionts which share at
 335 least one country in the distribution range, based on AntWiki (2019)] and the number of hits
 336 for their binomial name (and common synonyms) on *Google Scholar* as a proxy for sample
 337 effort for every ant species. Next we correlated total symbiont diversity with the host
 338 predictors described above, while correcting for the phylogenetic relatedness of the different
 339 ant species. The phylogenetic relatedness of host species should be accounted for as closely
 340 related host species cannot be treated as independent observations. For that reason, we
 341 modelled a phylogenetic generalized least-squares regression (PGLS) with total number of
 342 symbionts per ant species as the dependent variable. A PGLS model incorporates a
 343 phylogenetic variance–covariance matrix in its error structure. We used the variance–
 344 covariance matrix based on the pairwise node counts retrieved from the phylogenetic tree of
 345 European ants by Arnan *et al.* (2017). The phylogenetic relatedness of 108 out of the 181 ant
 346 species found in our data set was determined in this tree (see Appendix S3). From this subset
 347 of 108 species, we were able to find values for all predictors for 96 species. Consequently, the

348 PGLS model was based on these 96 ant species (Appendix S4). This data set reduction is
 349 acceptable as these 96 ant species interact with 620 symbionts and cover 86.5% of the
 350 recorded interactions in the host species–symbiont data set. The phylogenetic covariance
 351 matrix was multiplied by Pagel’s λ , a widely used parameter that scales and corrects for the
 352 expected degree of phylogenetic covariance (Pagel, 1999). This multiplier spans from 0,
 353 which corresponds to the complete absence of a phylogenetic signal in the residuals of the
 354 model (the model is then similar to a regular GLM with an ordinary least-squares error
 355 structure) to 1, when the covariance of the model residuals follows a Brownian motion model
 356 of evolution (Pagel, 1999; Freckleton, Harvey, & Pagel, 2002). The λ parameter
 357 characterizing the phylogenetic signal was estimated through maximum likelihood estimation
 358 within the PGLS model. We analysed this model using the *pgls* function embedded in the R
 359 3.5.1-package ‘caper’. We transformed the variables to meet the normality assumptions of the
 360 residuals. Number of symbionts was square–root transformed, the predictors colony size, the
 361 number of sympatric ants and *Google Scholar* hits ln-transformed, and the distribution range
 362 was square-root transformed. Finally, all continuous predictors were scaled to unit variance.
 363 In addition to this analysis on the drivers of total symbiont diversity, we ran similar PGLS
 364 models with subsets of symbiont species richness as dependent variables (overall number of
 365 myrmecophiles, trophobionts, social parasites and helminths, separately) and predictors of the
 366 subset of ant species that engage with these symbionts as predictors. Identical transformations
 367 of predictors and subsets of symbiont richness were applied as in the analysis on total
 368 symbiont richness described above. Diversity of fungi was not regressed against ant predictors
 369 in separate PGLS models as the number of host ants is relatively low in these groups.
 370 Models were ranked per analysis with the *dredge* function in the ‘MuMIn’ R-package
 371 according to their corrected Akaike Information Criterion (AICc) value. We retained the best

models identified with $\Delta AICc < 2$. Significance levels of the predictors of the retained models were assessed using Wald χ^2 tests.

(5) Predictors of symbiont sharing in European ants

Studying the factors that facilitate or constrain the transmission of a symbiont from one host to another is pivotal to understanding the ecological and co-evolutionary processes in host–symbiont networks (Okuyama & Holland, 2008; Pilosof *et al.*, 2013). For example, symbionts that are only shared by closely related hosts may indicate host–symbiont co-evolution in the network (Guimarães, Jordano & Thompson, 2011). Predictors were similar to the previous analysis and encompassed traits of the ant species and variables related to their habitat and distribution, including colony size, worker size, nest type, phylogeny as a proxy for trait similarity, habitat, distribution range, biogeographic region and allopatric distribution. We used multiple regression on distance matrices (MRM), an extension of partial Mantel analysis, to test the association between different distance matrices (Lichstein, 2007). The symbiont dissimilarity matrix contained the pairwise Jaccard distances between each pair of host ants based on the presence–absence data of the symbionts they supported. This matrix was regressed against multiple distance/dissimilarity matrices giving dissimilarities in the aforementioned predictors. Worker size difference was the absolute difference for this trait between every pair of ant species. The pairwise differences in colony size were ln-transformed. For the allopatric distribution matrix, we first calculated the degree of overlap in distribution (sympatry) between each pair of species (ln-transformed number of countries in the distribution range that are shared between each pair of ant species). We standardized this matrix between 0 and 1 and subtracted it from 1 to obtain a dissimilarity matrix. Entries in the distance matrices of habitat, nest type and biogeographic region were coded 0 when the pair of ants occupy the same habitat, nest type or biogeographic region, respectively, and 1 when

the pair of ants show differences in these variables. The phylogenetic distances were the pairwise node counts. Again, we focused our analysis on the subset of 96 ants for which the phylogenetic relationship was resolved by Arnan *et al.* (2017) and for which we possessed values for all predictors. We also included a matrix of sampling effort in which we pairwise multiplied the (ln+1)-transformed hits on *Google Scholar* of one ant species with the (ln+1)-transformed hits on *Google Scholar* of another ant species. All matrices were standardized between 0 and 1 and MRM analyses were conducted in the R package ‘ecodist’ using the MRM function. Significance of the predictor matrices was tested using 9999 permutations. We removed non-significant predictors, and reran the MRM analysis until all predictors were significant (Martiny *et al.*, 2011). The relative importance of the significant predictor matrices was calculated with the lmg metric, which uses unweighted averages of sequential R^2 of different orderings of the model predictors. The calculation and visualization of the lmg metrics was conducted with the R-package ‘relaimpo’.

We conducted similar MRM analyses on subsets of the symbiont community, where the response variable was the dissimilarity (pairwise Jaccard indices) in the set of myrmecophiles, trophobionts, social parasites, helminths and parasitic fungi of the host ants, respectively.

All statistical tests were conducted in R 3.5.2.

III. RESULTS

(1) Ant–symbiont networks display a diversity of species interactions

We identified 722 obligate ant symbionts interacting with 181 ant species in Europe (Appendix S1). The references we used to compile the host–symbiont interaction matrix are listed per symbiont species in Appendix S2. Myrmecophiles ($N = 535$) outnumbered the four other types of ant symbionts ($N = 80$, $N = 71$, $N = 22$ and $N = 14$ for trophobionts, social parasites, parasitic helminths and fungi, respectively). One fungus was mutualistic, the other

13 species were parasitic. Within the group of myrmecophiles, beetles and mites were the most species-rich groups (Fig. 1). The subdivision of myrmecophiles into parasitoids ($N = 33$), unspecialized myrmecophiles ($N = 413$) and specialized myrmecophiles ($N = 89$) can be found in Appendix S1. The hosts of 75 symbionts were not identified at the species level in the literature record. The distribution of the number of host species per symbiont was right-skewed (mean = 3.58, median = 2). The highest frequency (39%) of symbionts interacted with one host species and a maximum number of 34 host species was documented in the myrmecophilous silverfish *Proateturina pseudolepisma*.

European ant genera contained a highly variable number of species (proportional to the genus font size in Fig. 2; Fig. S1), ranging from 1 to 82, in the genera that interact with symbionts. Generally, the species-rich ant genera, such as the Formicinae genera *Lasius*, *Formica* and *Camponotus* and the Myrmicinae genera *Messor* and *Myrmica* attracted a higher diversity of all five types of symbionts (Pearson's product-moment correlation: $r = 0.59$, $P < 0.001$, d.f. = 30; Fig. S1). A notable exception was the European ant genus *Temnothorax*, which contained the most species while supporting a relatively moderate number of symbionts. Myrmecophiles were the dominant group in most ant genera. Trophobionts were generally the second most diverse group, but were absent or nearly absent in some genera such as *Monomorium*, *Aphaenogaster*, *Leptothorax*, *Messor*, *Temnothorax* and *Cataglyphis*. Social parasites contributed slightly to total symbiont diversity in most ant genera, but were very diverse in the ant genera *Temnothorax*, *Tetramorium* and *Leptothorax*. Helminths and fungi represent a minor fraction of the symbiont fauna in almost all ant genera. However, fungi are well-represented in *Myrmica*. Ant genera shared many symbionts with other genera (78.9% of the hosted community on average), belonging to the same or different ant subfamilies (connecting lines in Fig. 2). *Temnothorax*, *Leptothorax* and *Messor* are characterized by a relatively high number of unique symbionts (see relatively large inner circles in Fig. 2).

(2) Host specificity in different symbiont types

After controlling for sampling effort, symbiont groups had significantly different host ranges (Kruskal-Wallis chi-squared = 44.97, d.f. = 6, $P < 0.001$; Fig. 3). Social parasites had the lowest number of host species (Fig. 3, *post-hoc* comparisons indicated with letter code). Other symbiont types interacted with a higher number of ants, but we did not detect significant differences among these other types (Fig. 3). Host range of symbiont groups without controlling for sampling effort can be found in Fig. S2.

The average taxonomic distance between host species targeted by a symbiont was significantly different among symbiont groups (Kruskal-Wallis chi-squared = 83.37, d.f. = 6, $P < 0.001$; Fig. S3). The approach where we omitted the symbionts with one host species (Fig. S3) and that where we assigned a taxonomic distance of 1 to these species (Fig. S4) provided similar results (relative differences among symbiont groups are very similar in Figs S3 and S4). Likewise, we found significant differences in average phylogenetic distance between host species across the symbiont groups (GLM, $F = 17.27$, d.f. = 6, $P < 0.001$; Fig. 4). Patterns in the average host phylogenetic distance of the different symbiont groups closely matched those of the average host taxonomic distance. Host species of social parasites were very closely related to each other (Figs 4 and S3). Parasitic fungi also exploited related host species. Helminths and myrmecophiles interacted with hosts that are much more unrelated on average. We did not find differences in host relatedness among myrmecophilous parasitoids, unspecialized and specialized myrmecophiles (Figs 4 and S3). Ants interacting with trophobionts showed the lowest relatedness (Fig. 4).

(3) Predictors of symbiont diversity in European ants

471 The number of symbionts is highly variable in ant species. Here we report the host drivers that
 472 affect total symbiont richness and diversity of four subsets of ant symbionts (myrmecophiles,
 473 trophobionts, social parasites and helminths). Total symbiont diversity was clearly positively
 474 correlated with colony size. This factor was highly significant (PGLS, $P < 0.001$) in the four
 475 top-ranking models (Table 2, Fig. 5). Habitat and distribution range of the host were also
 476 incorporated in most of the top-ranking models. In these models, symbiont richness increased
 477 with the host distribution range and was highest in eurytopic habitats (PGLS, $P < 0.001$). As
 478 expected, sample effort has a major effect on the reported total symbiont diversity and the
 479 other subsets of symbiont diversity. Symbiont interactions were highest in ants that are
 480 intensively studied. We controlled for sample effort by including the proxy $(\ln+1)$ -
 481 transformed *Google Scholar* hits in our models. Myrmecophile richness was also positively
 482 affected by colony size (PGLS, $P < 0.001$ in the five top-ranking models, Table 2, Fig. 5),
 483 distribution and eurytopic habitat. Myrmecophiles were more abundant in larger ants (PGLS,
 484 $P < 0.05$ in the retained models). Trophobiont diversity was positively correlated with
 485 sampling effort and an eurytopic habitat in most models (Fig. 5, Table 2). Trophobiont
 486 diversity was in some models also higher in ant species with a higher distribution. There were
 487 no predictors consistently present in the top-ranking models explaining social parasite species
 488 richness (Table 2) Helminth diversity was higher in eurytopic ant species, but no other
 489 predictors were consistently retained in the top models (PGLS, $P < 0.001$, Table 2).
 490 We found a phylogenetic signal in the predictors of the PGLS models ($\Delta AICc < 2$) with total
 491 symbiont richness as dependent variable (Pagel λ ranged from 0.41 to 0.54 Table 2). The
 492 residuals of the models showed different degrees of phylogenetic covariance. The largest
 493 phylogenetic signal was found in the models with social parasites (Pagel λ ranged from 0.83
 494 to 1.00) and helminths (Pagel λ ranged from 0.85 to 0.94) as dependent variable. By contrast,

phylogenetic relatedness of the hosts did not explain additional variation (Pagel's $\lambda = 0$) in most retained models with myrmecophiles as dependent variable (Table 2).

(4) Predictors of symbiont sharing in European ants

The dissimilarity matrices of the host predictors (indicated with Δ in Fig. 6) were positively correlated with dissimilarity in symbiont composition (Fig. 6). This shows that ant species with a higher similarity in these predictors display a higher similarity in symbiont composition. The most important predictors of similarity in ant symbiont communities in the European ant data set were phylogenetic relatedness of the ant hosts (MRM, $\text{lmg} = 0.40$, $P < 0.001$) and similarity in biogeographic region (MRM, proportional contribution to the total $r^2 = \text{lmg} = 0.35$, $P < 0.001$) (Fig. 6). Similarities in worker size ($\text{lmg} = 0.07$, $P = 0.01$), colony size ($\text{lmg} = 0.04$, $P = 0.02$) and habitat ($\text{lmg} = 0.01$; $P = 0.03$) also facilitated the sharing of symbionts (Fig. 6). Better studied ant pairs shared more similar symbiont communities (MRM, $\text{lmg} = 0.13$, $P < 0.001$). Well-studied ant pairs also shared more trophobionts, myrmecophiles and parasitic fungi (MRM-analyses, lmg ranging from 0.14-0.60, all $P < 0.05$, Fig. 6). Trophobiont sharing was also positively correlated with phylogenetic relatedness and similarity in biogeographic regions of the ant hosts (MRM, $\text{lmg} = 0.22$, $P < 0.001$ and $\text{lmg} = 0.13$, $P < 0.001$, respectively). The similarities in social parasite communities was largely explained by phylogenetic relatedness (MRM, $\text{lmg} = 0.76$, $P < 0.001$). Similarities in biogeography (MRM, $\text{lmg} = 0.04$, $P = 0.04$) explained additional variation in the sharing of social parasites. Helminth sharing was also strongly facilitated in phylogenetically related ant hosts (MRM, $\text{lmg} = 0.71$, $P < 0.001$). Interestingly, phylogenetic relatedness of the hosts did not promote the sharing of myrmecophiles. The similarity of myrmecophile communities between ant hosts was mainly driven by living in a similar biogeographic region (MRM, lmg

= 0.68, $P < 0.001$), Lastly, ants in the same biogeographic region tend to share more parasitic fungi (MRM, $\text{Im}g = 0.44$, $P < 0.01$) (Fig. 6).

IV. DISCUSSION

Understanding community composition and stability is an important challenge in ecology. Network analysis has approached this challenge, using community structure and species interactions as fundamental building blocks. Yet, studies that explain the detailed topology of large-scale ecological networks encompassing a diversity of interaction types are limited. We here provide a complete tally of the distribution of ant symbiont groups over European ants and compare host specificity, symbiont richness, host switching and its drivers for different ant symbiont groups.

(1) Characterization of the European ant–symbiont network

It is widely acknowledged that the group of obligate ant symbionts is hyperdiverse (Wasmann, 1894; Kistner, 1979, 1982; Hölldobler & Wilson, 1990; Rettenmeyer *et al.*, 2010), although exact species numbers at a regional scale are lacking. Rough estimates of the global diversity of parasites living in ant nests reach 10,000 to 20,000 species (Thomas, Schönrogge & Elmes, 2005), which is higher than mammal and bird diversity. We here identified 722 symbionts distributed over 181 ant species in Europe. The majority of these symbionts were classified as myrmecophiles, which are commensalistic-to-parasitic arthropods mostly living inside the ant nest (Kronauer & Pierce, 2011; Parmentier *et al.*, 2016a; Parmentier, 2020). Beetle and mite communities were the most diverse groups. In other regions, beetles and mites also outnumber other myrmecophilous arthropod groups (Kistner, 1982; Hölldobler & Wilson, 1990). Social parasites and mutualistic trophobionts are medium-sized groups; endoparasitic helminths and parasitic fungi are relatively species-poor, but understudied.

Mutualistic ant symbionts are thus clearly overshadowed by the diversity of commensalistic and parasitic ant symbionts in Europe. Species-rich ant genera and subfamilies generally supported higher numbers of ant symbionts. Host–symbiont networks are characterized by an asymmetrical organization of interactions with host-specific symbionts and symbionts that interact with multiple host taxa (Guimarães *et al.*, 2006). Overall, a large proportion of the symbionts were shared among heterogeneric ant species (Fig. 2). Some ant genera interacted with a relatively low number of symbionts, but most of their symbionts were not found in association with other ant genera. The highly specific group of social parasites was much more represented in the symbiont community of these hosts. In addition, the group of trophobionts is diverse in some ant genera, but is marginal or even absent in others (Fig. 2). The distribution of trophobiont interactions across the ant genera echoed the dietary preferences of the host. The diet of genera such as *Formica*, *Lasius* and *Myrmica* consists of a significant proportion of honeydew, whereas genera with few or no trophobiotic interactions are known to be predatory (e.g. *Temnothorax*, *Cataglyphis*) or granivorous (e.g. *Messor*) (Seifert, 2007). Mutualistic interactions are much more diverse in tropical systems than those observed in the European network and include ant-cultivated plants, ant-defended plants and ant-cultivated fungi (Rico-Gray & Oliveira, 2007). The uneven distribution of the five types of symbionts among the European ant genera suggests that some ant lineages are more predisposed to associate with particular types of symbionts. Ant–host associations are thus shaped by deep evolutionary processes as determined by biotic and environmental drivers of speciation and extinction (Aguilée *et al.*, 2018).

(2) Host specificity in different symbiont types

Host specificity is a key feature of host–symbiont networks, and is moulded by the ecological and evolutionary interactions between the host and symbiont (Poulin & Mouillot, 2003).

569 Patterns in host specificity have been studied in a wide range of host–symbiont systems.
570 Generally, parasites are thought to have a tendency to evolve to extreme host specialization as
571 they need complex adaptations to bypass host defences (Kawakita *et al.*, 2010). The drivers
572 favouring host specificity in mutualist partners are far less understood and both low and high
573 degrees of specificity are widespread (e.g. plant–seed dispersers and fig–fig wasps,
574 respectively). Ant symbioses are ideal to unravel patterns in host specificity. They occupy the
575 complete mutualism–parasitism continuum and allow comparison of host specificity in
576 different types of symbionts. We here demonstrate that average host range in European ant
577 symbionts was much broader than previously assessed in a study on host specificity of
578 myrmecophiles at a global scale (Glasier *et al.*, 2018) which found that obligate ant symbionts
579 occurred on average with *ca.* 1.20 host species. We, however, found that European symbionts
580 were reported with three times this number of host species (3.58) on average. The much lower
581 number of detected hosts in Glasier *et al.* (2018) is probably the result of their searching
582 method. They did not include data from faunistic notes, grey literature and books, which
583 report the majority of interactions between ants and their symbionts. Moreover, the symbiont
584 fauna, let alone the range of their interactions, is poorly documented outside Europe, which
585 makes hard predictions at a global scale unreliable (Parmentier, 2020). Ant symbionts were
586 extremely variable over the host-ant range. After controlling for sampling effort, social
587 parasites clearly targeted the lowest number of host species, which is in line with expectations
588 as they are the most specialized group of parasites (Buschinger, 2009). Apart from the number
589 of hosts, the relatedness of host species is also a vital aspect of host specificity. It is well
590 described that social parasites colonize nests of related hosts (Emery’s rule; Buschinger,
591 2009). However, this has not been compared with other types of symbiont groups. We showed
592 that the hosts of social parasites were clearly the most related of all symbiont types. The hosts

of myrmecophiles, parasitic fungi and helminths showed moderate to poor relatedness on average. Trophobionts were associated with the most distantly related ant species. There is a large body of literature that explains the constraints of host switching in social insect symbionts. Generally, it is thought that specialized myrmecophiles and social parasites rely on chemical deception, by mimicking the colony recognition cues or some key pheromones (overview in Parmentier, Dekoninck & Wenseleers, 2017). They are completely integrated into the host colony and are treated as a true colony member. Because of this strict mimicking of the host's communication system, they are not able to colonize unrelated host species. Unspecialized myrmecophiles are typically poorly integrated into the colony, but host switching is more common in this group. This is facilitated by the use of general defensive chemicals, chemical insignificance or behavioural strategies (Stoeffler, Tolasch & Steidle, 2011; Parmentier *et al.*, 2017, 2018). Consequently, we predicted that specialized myrmecophiles would display much higher degrees of host specificity than unspecialized myrmecophiles. Surprisingly, we did not find differences in the number of host species and host relatedness among unspecialized myrmecophiles, specialized myrmecophiles and parasitoids. A limitation of the present study is that we do not have information on the relative importance of the recorded host species for a symbiont. It is likely that some of the listed host species of specialized symbionts rarely act as hosts, resulting in an overestimation of the actual species range and host species relatedness of symbionts. Additionally, the biology of most myrmecophiles is poorly known, which makes a functional grouping according to specialization challenging and open for refinement. The processes which make trophobionts attractive to one host, but not to another are hitherto unexplored. Likewise, the mechanisms of host switching and the factors that facilitate or constrain host switching in endoparasitic helminths and fungi are unknown.

(3) Predictors of symbiont diversity in European ants

Associations between ant hosts and their symbionts are not random and are structured according to both ecological and evolutionary factors that act at different spatiotemporal scales. From the perspective of ant symbionts, ant nests can be conceptualized as habitat islands. Ant species with larger ant nests interact with more symbiont species. Nest size of ant species has been repeatedly hypothesized as an important driver of ant symbiont diversity (Hughes, Pierce & Boomsma, 2008; Kronauer & Pierce, 2011), and here was formally tested for the first time. Previous studies across very different host–symbiont systems [e.g. ectoparasites of fishes (Guégan *et al.*, 1992), parasites of hoofed mammals (Ezenwa *et al.*, 2006), parasites of carnivores (Lindenfors *et al.*, 2007), feather mites of finches (Villa *et al.*, 2013)] identified the size of the host species as one of the key factors in determining symbiont species richness (Kamiya *et al.*, 2014). This positive association results from the fact that larger host species provide more niches and are less ephemeral (Lindenfors *et al.*, 2007). Analogously, ant species with larger nests provide more space to allow larger population sizes, thereby reducing the extinction risk of symbionts (*cf.* island theory; Macarthur & Wilson, 1967). In addition, ant species with larger nests provide a higher diversity of microhabitats, including refuge areas that eventually facilitate species coexistence (Barabás *et al.*, 2018). Larger ant nests are also expected to be more persistent (Kaspari & Vargo, 1995). The colony size of ant species is thus a strong local driver of total symbiont richness, and myrmecophile richness in particular.

Total symbiont diversity, myrmecophile diversity, helminth and trophobiont diversity are additionally determined by more regional ecological factors like range size and niche width of the host ants. In that respect, eurytopic ants, such as *Lasius niger* and *Myrmica rubra* that can live in a wide variety of habitats including urban regions, hosted more symbionts, myrmecophiles, trophobionts and helminths. An effect of both distribution and habitat reflects

that more symbionts occur in widely distributed ant species with high densities. Symbionts associated with widely distributed ants are less prone to extinction as predicted by life-history theory and metacommunity ecology (Nosil, 2002; Leibold *et al.*, 2004).

Host density has widely been demonstrated as a key factor explaining parasite species richness (Lindenfors *et al.*, 2007). Interestingly, we found that species with larger workers engaged with more myrmecophiles than ant species with small workers. This pattern in myrmecophile diversity was previously hypothesized based on experimental work, showing that the survival of three myrmecophilous beetles gradually increased in laboratory nests of larger ant species (Parmentier, Dekoninck & Wenseleers, 2016b). This positive relationship between ant size and myrmecophile diversity suggests that species with small workers detect, attack and deter myrmecophiles more easily and efficiently.

Sampling effects appear highly relevant in most models of symbiont diversity among different hosts. Trophobiotic and more specialized parasitic interactions as seen in the group of helminths and social parasites are more determined by evolutionary drivers than myrmecophiles. Indeed, affinity with their hosts is strongly shaped by the phylogeny of the host. The effect of phylogeny is echoed in the high Pagel's lambda values of the corresponding PGLS models, implying that much of the residual variation in trophobiont and especially helminth and social parasite richness could be explained by the phylogeny of the hosts. This strong phylogenetic driver for social parasite richness is in line with our previous results that social parasites mainly target closely related ant species (referred to as Emery's rule; Buschinger, 2009) belonging to a small number of ant genera. Symbiont network structure thus shifts from more neutral ecological drivers related to regional species abundance to co-evolutionary drivers related to ancestry. The uniqueness and tightness of species interactions are known to be both a driver and consequence of co-evolutionary dynamics. Interestingly, we here show that these evolutionary drivers overrule any ecological

one in the most specialized interactions (social parasites), hence demonstrating the integrated nature of symbiont network formation according to the prevailing interaction strengths. Other predictors, which were not considered in the analyses, may also positively affect the diversity of symbionts. The availability of larvae and pupae rather than colony size may be more important for parasitoid species (Rocha *et al.*, 2020). The tolerance level of ant species to intruders and the colonial organization (either single nests or multiple connected nests, either a single queen or multiple queens) are also possible determinants of diversity, but sufficient information is lacking to test these hypotheses.

Ant–symbiont networks are unique in the sense that the host-associated network that is studied covers a wide array of interactions, from putatively mutualist to strictly antagonistic. We found that evolutionary processes are pivotal in networks of the most specialized ant symbionts (social parasites), whereas less-specialized networks, as found in the group of myrmecophiles, were mostly determined by ecological factors. The same pattern was found in other symbiont systems. Studies on specialized host–parasite networks equally point at the dominance of evolutionary drivers (phylogeny and biogeography) of these associations (Feliu *et al.*, 1997; Rosas-Valdez & de Pérez-Ponce de León, 2011), while less-obligatory animal parasitic (Nunn *et al.*, 2003; Ezenwa *et al.*, 2006; Lindenfors *et al.*, 2007; Nava & Guglielmone, 2013) or plant mutualistic interactions (Sanders, 2003; Wagner, Mendieta-Leiva, & Zotz, 2015) are more affected by ecological factors related to distribution and abundance patterns that enhance contact and hence transmission of their diversity and host-association patterns.

(4) Predictors of symbiont sharing in European ants

We hypothesized that the shared evolutionary history of related ant species would promote the sharing of similar symbiont communities. A positive correlation between phylogenetic

relatedness of the hosts and symbiont sharing was demonstrated in previous studies on orchid mycorrhiza (Jacquemyn *et al.*, 2011) and bat viruses (Luis *et al.*, 2015), but no such relationship was found in arbuscular mycorrhiza (Veresoglou & Rillig, 2014) and primate parasites (Cooper *et al.*, 2012). Consistent with our prediction, we found that the main factor that promoted symbiont sharing in European ants was the relatedness of the hosts. It indicates that many symbionts pass more easily to related host species. As related ant species employ nearly identical defence structures (nestmate recognition cues, physiological and behavioural defences), it enables symbionts, especially specialized parasites, to bypass the host defence systems of related hosts. Another key factor that may facilitate the cross-species transmission of symbionts is the overlap in geographical distribution of the hosts (*cf.* bat viruses in Luis *et al.*, 2015). We showed that ant species living in the same biogeographical region possessed more similar symbiont communities. This suggests that both the spatial overlap and similarity in climatic conditions facilitate the sharing of symbionts. Sampling effort also considerably explained the sharing of symbionts. More shared symbionts were reported in well-studied pairs of species. Focusing on the different subsets of ant symbionts, we found that the sharing of trophobionts between host ant species was mainly determined by biogeography and phylogenetic relatedness. Phylogenetic relatedness of the hosts was the most important driver explaining the sharing of helminths and social parasites. The strong phylogenetic effect on the sharing of social parasites is directly linked to the very low taxonomic/phylogenetic distance between their hosts. Social parasites target a very narrow range of host species by hijacking their communication system. This exploitation of host cues is facilitated by immediate common ancestry (Buschinger, 2009). The biology of most helminths is unknown, but probably immune evasion is only possible in related host species. Myrmecophiles and parasitic fungi were more similar in ant species residing in the same biogeographical region. Climatic conditions have a strong effect on the distribution of different groups of

myrmecophiles. One example is the large group of myrmecophilous silverfish which are mainly confined to ants living in the Mediterranean region (Molero-Baltanás *et al.* 2017; Appendix S1). Interestingly, host switching of myrmecophiles and parasitic fungi was not positively correlated with host relatedness (*cf.* Cooper *et al.*, 2012; Veresoglou & Rillig, 2014).

V. OUTSTANDING QUESTIONS

Merging different interaction types into one ecological network framework is a key challenge in ecology (Fontaine *et al.*, 2011). Diverse host–symbiont communities provide an opportunity to test the relative contributions of ecology and evolution to network assembly. For example, our study on ant–symbiont networks revealed different roles of ecological and evolutionary processes depending on the type of symbiosis. Our insights may provide a basis for theory development and across-ecosystem comparisons (e.g. plant- and coral-based networks) and synthesis.

We lack theory on how the architecture and the interaction signs and sizes jointly affect the stability and productivity of these diverse networks, much in contrast to trophic or mutualistic networks. The relative ease with which one can manipulate ant–symbiont communities makes them suited as empirical systems to test theory.

Host–symbiont networks offer an opportunity to understand both ecological and evolutionary processes behind community assembly, from meso- to macro-ecological scales (see Vellend, 2016). More specifically, as hosts occur spatially structured at these scales, it remains an open question how these assembly processes are determined by ecological and evolutionary limitation of dispersal. One key question is whether and how symbionts are dispersing: to what degree is horizontal transfer and subsequent symbiont sharing across hosts a facilitator of symbiont community assembly, and to which degree is vertical transfer, i.e. co-dispersal of

743 symbiont and host, established across the antagonism–mutualism gradient of host–symbiont
 744 networks? Mutualistic plant mycorrhizal fungi and plant diaspores, for instance, are passively
 745 co-dispersed by birds (Correia *et al.*, 2019). Are similar processes equally prevalent in ant–
 746 symbiont interactions, for instance are symbionts transported by their host during colony
 747 relocation (Parmentier, 2019)?
 748 Empirical data demonstrate that different host ants coexist regionally. The stabilizing and
 749 equalizing mechanisms that underpin such coexistence are at present unknown. How do the
 750 complex symbiont networks in which these hosts are embedded contribute to such
 751 mechanisms? Addressing such questions with new analytical tools in coexistence research
 752 (Saavedra *et al.*, 2017; Spaak & De Laender, 2020) could advance our basic understanding of
 753 how a variety of direct and indirect interactions affect coexistence among hosts. For example,
 754 do symbionts induce indirect interactions among ant hosts? Do ant density-dependent
 755 interactions between symbionts (Parmentier *et al.*, 2018) represent a stabilizing higher-order
 756 interaction (Grilli *et al.*, 2017)?
 757 Insights from this review are restricted and applicable to networks as characterized at the
 758 species level, thereby neglecting any intraspecific variation. Following the relevance of
 759 ecological and evolutionary determinants, the question remains open as to what degree co-
 760 evolutionary dynamics between hosts and their symbiont community occur. As strong
 761 selection may act on ant symbionts to bypass host colony defence, cryptic speciation in ant
 762 symbionts is expected to be high (Schönrogge *et al.*, 2002; Zagaja & Staniec, 2015; von
 763 Beeren, Maruyama & Kronauer, 2015). Symbiont populations may be adapted to an
 764 individual host population as was demonstrated in the ant-parasitic syrphid fly *Microdon* and
 765 the butterfly *Phengaris* (Elmes *et al.*, 1999; Tartally *et al.*, 2019). Ultimately, population
 766 divergence may result in cryptic symbiont species each targeting a different host species.

At a higher phylogenetic level, other hymenopteran and insect lineages (Isoptera) provide similar niches to nest symbionts. None of the listed ant symbionts are shared with wasps, solitary and eusocial bees and termites (note that the latter two groups are poorly represented in Europe). Apparently only facultative symbionts (e.g. *Porcellio scaber* in wasp and bee nests for instance) or very generalist entomopathogens such as *Beauveria bassiana* are able to spread across different social insect lineages, but more study is needed to understand the drivers of host–symbiont divergence at these deep phylogenetic levels.

VI. CONCLUSIONS

(1) Ant–symbiont networks are particularly interesting to study large-scale patterns and drivers in host–symbiont network topology and symbiont richness as they are extremely diverse and cover the entire mutualism–antagonism continuum. We assembled a complete network of ant–symbiont interactions in Europe and studied the drivers of host specificity, symbiont richness and symbiont sharing in the different interaction sub-networks.

(2) We identified 722 ant macrosymbionts which we categorized in five types: (1) myrmecophiles – commensalistic and parasitic arthropods ($N = 535$); (2) trophobionts – mutualistic aphids, scale insects, plant hoppers and mutualistic Lepidoptera ($N = 80$); (3) social parasites – parasitic ant species ($N = 71$); (4) parasitic helminths ($N = 22$); (5) fungi – parasitic ($N = 13$) and mutualistic ($N = 1$).

(3) The different types of ant symbionts significantly varied in host specificity. Apart from quantitative differences in host range, we also found clear differences in the average taxonomic/phylogenetic relatedness of the targeted host species for the different types of ant symbionts. The most species rich and best-studied ant genera generally supported the largest number of symbionts, but the different types of symbionts were unevenly distributed across ant genera.

(4) We revealed that the ecological and evolutionary factors which drive symbiont species richness may shift depending on the type of symbiosis. Myrmecophile species richness is mainly determined by ecological drivers, such as colony size, host range and niche width of the host. By contrast, species richness of social parasites is strongly determined by the evolutionary factor host phylogeny.

(5) Ants living in the same biogeographic region shared more symbionts. The sharing of trophobionts, helminths and social parasites, in particular, was also strongly facilitated in phylogenetically related hosts.

VII. ACKNOWLEDGEMENTS

This work was supported by Bijzonder Onderzoekfonds Ugent (BOF17/PDO/084 to T.P.) and FWO (grant nr. 1203020N to T.P.). D.B. and F.d.L. are supported by the FWO research network EVENET (W0.003.16N). We thank the anonymous reviewers for their helpful and constructive comments.

VIII. REFERENCES

References marked with an asterisk are cited only within the supporting information.

* AGRAIN, F.A., BUFFINGTON, M.L., CHABOO, C.S., CHAMORRO, M.L. & SCHÖLLER, M. (2015). Leaf beetles are ant-nest beetles: The curious life of the juvenile stages of case-bearers (Coleoptera, Chrysomelidae, Cryptocephalinae). *ZooKeys* **2015**, 133–164.

AGUILÉE, R., GASCUEL, F., LAMBERT, A. & FERRIERE, R. (2018). Clade diversification dynamics and the biotic and abiotic controls of speciation and extinction rates. *Nature Communications* **9**, 3013.

* AKYÜREK, B., ZEYBEKOĞLU, Ü., GÖRÜR, G. & KARAVIN, M. (2016). Reported aphid (Hemiptera: Aphidoidea) and ant (Hymenoptera: Formicidae) species associations from

- 817 Samsun province. *Journal of the Entomological Research Society* **18**, 97–106.
- 818 * ALBRECHT, A.C. (2017). Illustrated identification guide to the Nordic aphids feeding on
819 Conifers (Pinophyta) (Insecta, Hemiptera, Sternorrhyncha, Aphidomorpha. *European*
820 *Journal of Taxonomy* **338**, 1–160.
- 821 ALLESINA, S. & TANG, S. (2015). The stability–complexity relationship at age 40: a random
822 matrix perspective. *Population Ecology* **57**, 63–75.
- 823 * ÁLVAREZ, M., MUNGUIRA, M.L. & MARTÍNEZ-IBÁÑEZ, M.D. (2012). Nuevos datos y
824 recopilación de las relaciones entre Lycaenidae y Formicidae en la Península Ibérica
825 (Lepidoptera: Lycaenidae; Hymenoptera: Formicidae). *SHILAP Revista de*
826 *Lepidopterologia* **40**, 45–59.
- 827 ANTWIKI (2019). AntWiki. [Http://www.antwiki.org/wiki/](http://www.antwiki.org/wiki/) [accessed 20 July 2019].
- 828 ARNAN, X., CERDA, X. & RETANA, J. (2017). Relationships among taxonomic, functional, and
829 phylogenetic ant diversity across the biogeographic regions of Europe. *Ecography* **40**,
830 448–457.
- 831 * ASSING, V. (1999). A revision of the genus *Piochardia* Heyden, 1870 (Insecta: Coleoptera:
832 Staphylinidae: Aleocharinae). *Annalen des Naturhistorischen Museums in Wien. Serie B*
833 *für Botanik und Zoologie* **101**, 277–301.
- 834 * ASSING, V. (2001). The first record of *Dinusa* Saulcy 1864 from Turkey (Coleoptera:
835 Staphylinidae, Aleocharinae). *Linzer biologische Beiträge* **33**, 187–190.
- 836 * ASSING, V. (2003). New species and records of *Eurysinius*, subgenus of *Astenus* Dejean, from
837 the Iberian Peninsula (Coleoptera: staphylinidae, Paederinae). *Linzer biologische Beiträge*
838 **35**, 693–700.
- 839 * ASSING, V. (2005). Two new species and new records of Staphylinidae from the Greek island

- 840 Lesbos (Insecta: Coleoptera). *Linzer biologische Beiträge* **37**, 1035–1046.
- 841 * ASSING, V. (2013a). On the Staphylinidae (Coleoptera) of Crete, Greece. *Stuttgarter Beiträge*
842 *zur Naturkunde A* **6**, 83–102.
- 843 * ASSING, V. (2013b). On the Staphylinidae of Rhodes, Greece (Insecta : Coleoptera). *Linzer*
844 *biologische Beiträge* **45**, 1587–1613.
- 845 * ASSING, V. (2014). On the Staphylinidae of Sardinia, Italy (Insecta : Coleoptera). *Linzer*
846 *biologische Beiträge* **46**, 415–428.
- 847 * ASSING, V. (2015a). On the Staphylinidae (Coleoptera) of Crete II. Seven new species, a new
848 synonymy, and additional records. *Stuttgarter Beiträge zur Naturkunde A, Neue Serie* **8**,
849 95–112.
- 850 * ASSING, V. (2015b). On the Staphylinidae of the Greek island Samos (Coleoptera :
851 Staphylinidae). *Koleopterologische Rundschau* **85**, 81–102.
- 852 * ASSING, V. (2016). On the Staphylinidae of the Greek island Karpathos (Insecta: Coleoptera).
853 *Linzer biologische Beiträge* **48**, 225–263.
- 854 * BACH, C. & CÁRDENAS-TALAVERÓN, A. (1984). Cita para Andalucía de *Amorphocephalus*
855 *coronatus* (Germar, 1817). (Col. Brentidae). *Miscellània Zoològica* **8**, 287–288.
- 856 * BALL, S. & MORRIS, R. (2015). *Britain's Hoverflies: A Field Guide*. Princeton University
857 Press.
- 858 * BALOGH, J. (1938). Studien über die myrmecophilen Milben Ungarns, I. *Folia entomologica*
859 *hungarica* **3**, 106–109.
- 860 BARABÁS, G., D'ANDREA, R. & STUMP, S.M. (2018). Chesson's coexistence theory.
861 *Ecological Monographs* **88**, 277–303.

862 * BARONI-URBANI, C. (1966). Osservazioni diverse intorno al nomadismo dell'*Aphaenogaster*
863 *picena* Baroni con particolare riguardo all'orientamento (Hymenoptera, Formicidae).
864 *Insectes Sociaux* **13**, 69–86.

865 BASCOMPTE, J. & JORDANO, P. (2007). Plant-animal mutualistic networks: The architecture of
866 biodiversity. *Annual Review of Ecology, Evolution, and Systematics* **38**, 567–593.

867 BASCOMPTE, J., JORDANO, P. & OLESEN, J.M. (2006). Asymmetric coevolutionary networks
868 facilitate biodiversity maintenance. *Science* **312**, 431–433.

869 BASTOLLA, U., FORTUNA, M.A., PASCUAL-GARCIA, A., FERRERA, A., LUQUE, B. &
870 BASCOMPTE, J. (2009). The architecture of mutualistic networks minimizes competition
871 and increases biodiversity. *Nature* **458**, 1018–1020.

872 * BÁTHORI, F., PFLIEGLER, W.P., ZIMMERMAN, C.-U. & TARTALLY, A. (2017). Online image
873 databases as multi-purpose resources: discovery of a new host ant of *Rickia wasmannii*
874 Cavara (Ascomycota, Laboulbeniales) by screening AntWeb.org. *Journal of Hymenoptera*
875 *Research* **61**, 95-104.

876 * BAUMANN, J. (2018). Tiny mites on a great journey – a review on scutacarid mites as phoronts
877 and inquilines (Heterostigmatina, Pygmephoroidae, Scutacaridae). *Acarologia* **58**, 192-
878 251.

879 * BEKCHIEV, R., SIVILOV, O., KOSTOVA, R. & LAPEVA-GJONOVA, A. (2017). Distribution of
880 *Amorphocephala coronata* (Germar, 1817). (Coleoptera: Brentidae) in Bulgaria. *ZooNotes*
881 **104**, 1–3.

882 * BERG, M. (1995). De mierenpissebed, *Platyarthrus hoffmannseggii*, een mysterieuze gast in
883 mierennesten. *Natura* **92**, 62–65.

884 * BEROS, S., JONGEPIER, E., HAGEMEIERS, F. & FOITZIK, S. (2015) The parasite's long arm: A
885 tapeworm parasite induces behavioural changes in uninfected group members of its social

- 886 host. *Proceedings of the Royal Society B: Biological Sciences* **282**.
- 887 * BINAZZI, A. & SCHEURER, S. (2009). *Atlas of the honeydew producing conifer aphids of*
888 *Europe*. ARACNE editrice S.r.l.
- 889 * BLACKMAN, R.L. & EASTOP, V.F. (2017). Aphids on the World's Plants: An Identification
890 and Information Guide <http://www.aphidsonworldsplants.info/> [accessed June 2018]
- 891 BLÜTHGEN, N., MENZEL, F., HOVESTADT, T., FIALA, B. & BLÜTHGEN, N. (2007)
892 Specialization, constraints, and conflicting interests in mutualistic networks. *Current*
893 *Biology* **17**, 341–346.
- 894 * BOLAÑO, J.A.S. & VILLERO, J.M.B. (2010). Los Colévidos epigeos (Coleoptera, Leiodidae,
895 Cholevinae) de la Sierra de Tudia (Badajoz, Extremadura, España). *Boletín de la Sociedad*
896 *Entomológica Aragonesa (S.E.A.)* **47**, 357–361.
- 897 BOULAY, R., ARON, S., CERDÁ, X., DOUMS, C., GRAHAM, P., HEFETZ, A. & MONNIN, T.
898 (2017). Social life in arid environments: The case study of *Cataglyphis* ants. *Annual*
899 *Review of Entomology* **62**, 305–321.
- 900 * BOUYON, H. & PERREAU, M. (2015). *Anemadus strigosus* Kraatz, 1852 en Alsace, première
901 citation en France (Coleoptera Leiodidae Cholevinae Anemadini). *L'Entomologiste* **71**,
902 137–139.
- 903 * BRAČKO, G. (2010). First record of a slave-making ant *Myrmoxenus gordiagini* Ruzsky, 1902
904 (Hymenoptera: Formicidae) in Slovenia. *Acta Entomologica Slovenica* **18**, 137–142.
- 905 * BREEN, J. (1979). Aphids visited by *Formica lugubris* (Hymenoptera : Formicidae) including
906 eleven species new to Ireland. *Irish Naturalist Journal* **19**, 349–352.
- 907 * BREWITT, K., PIÑOL, J., WERNER, C., BEYSCHLAG, W., ESPADALER, X., PÉREZ HIDALGO, N.
908 & PLATNER, C. (2014). Evaluating the importance of trophobiosis in a Mediterranean ant

- 909 community: a stable isotope analysis. *Insectes Sociaux* **62**, 81–95.
- 910 * BRIAN, M.V., HIBBLE, J. & STRADLING, D. (1965). Ant pattern and density in a southern
911 English heath. *Journal of Animal Ecology* **34**, 545–555.
- 912 BRONSTEIN, J.L., ALARCÓN, R. & GEBER, M. (2006). The evolution of plant-insect
913 mutualisms. *New Phytologist* **172**, 412–428.
- 914 * BURAKOWSKI, B. & NEWTON A. (1992). The immature stages and bionomics of the
915 myrmecophile *Thoracophorus corticinus* Motschulsky, and placement of the genus.
916 *Annali del Museo Civico di Storia Naturale di Genova* **89**, 17–42.
- 917 * BUSCHINGER, A. (1989a). Evolution, speciation, and inbreeding in the parasitic ant genus
918 *Epimyrma* (Hymenoptera, Formicidae). *Journal of Evolutionary Biology* **2**, 265–283.
- 919 * BUSCHINGER, A. (1989b). Workerless *Epimyrma kraussei* Emery 1915, the first parasitic ant
920 of Crete. *Psyche: A Journal Of Entomology* **96**, 69–74.
- 921 BUSCHINGER, A. (2009). Social parasitism among ants : a review (Hymenoptera : Formicidae).
922 *Myrmecological News* **12**, 219–235.
- 923 * BUSCHINGER, A., EHRHARDT, W. & FISCHER, K. (1981). *Doronomyrmex pacis*, *Epimyrma*
924 *stumperi* und *E. goesswaldi* (Hym., Formicidae) neu für Frankreich. *Insectes Sociaux* **28**,
925 67–70.
- 926 * BUSCHINGER, A., EHRHARDT, W., FISCHER, K. & OFER, J. (1988). The slave-making ant genus
927 *Chalepoxenus* - review of literature, range, slave species. *Zoologische Jahrbücher,*
928 *Abteilung für Systematik, Ökologie und Geographie der Tiere.* **115**, 383–401.
- 929 * BUSCHINGER, A. & WINTER, U. (1985). Life history and male morphology of the workerless
930 parasitic ant *Epimyrma corsica* (Hymenoptera: Formicidae). *Entomologia Generalis* **10**,
931 65–75.

- 932 * BUSCHINGER, A., WINTER, U. & FABER, W. (1983). The Biology of *Myrmoxenus gordiagini*
 933 Ruzsky, a slave-making ant (Hymenoptera, Formicidae). *Psyche: A Journal Of*
 934 *Entomology* **90**, 335–342.
- 935 CAGNOLO, L. & TAVELLA, J. (2015). The network structure of myrmecophilic interactions.
 936 *Ecological Entomology* **40**, 553–561.
- 937 * CAMMAERTS, R., CAMMAERTS, M. & DETRAIN, C. (1989) Response of the myrmecophilous
 938 beetles *Edaphopausus favieri* (Carabidae, Paussinae) and *Dichillus minutus*
 939 (Tenebrionidae) to the trail of their host *Pheidole pallidula*. *Actes des Colloques Insectes*
 940 *Sociaux* **5**, 199–206.
- 941 * CARLES-TOLRÁ, M. (2006). *Microselia micropila* sp.n.: a new phorid species from Spain,
 942 ethology of *Microselia rivierae* Schmitz, 1934, and a key to the European species
 943 of *Microselia* Schmitz (Diptera, Phoridae). *Boletín de la Sociedad Entomológica*
 944 *Aragonesa* **39**, 103–109.
- 945 * CARTER, C.I. & MASLEN, N.R. (1982). *Conifer lachnids*. Forestry Commission Bulletin 58.
 946 HMSO, London.
- 947 * CONSTANTINESCU, I., IVAN, O., CĂLUGĂR, A. & MARKÓ, B. (2011). Mite Fauna of Ant Nests
 948 - Comparative study of mite fauna in the Argeş river basin (South Romania). *Travaux du*
 949 *Muséum National d'Histoire Naturelle 'Grigore Antipa'* **54**, 327–342.
- 950 COOPER, N., GRIFFIN, R., FRANZ, M., OMOTAYO, M. & NUNN, C.L. (2012). Phylogenetic host
 951 specificity and understanding parasite sharing in primates. *Ecology Letters* **15**, 1370–1377.
- 952 CORREIA, M. , HELENO, R. , DA, SILVA, L. P., COSTA, J. M. & RODRÍGUEZ-ECHEVERRÍA, S.
 953 (2019), First evidence for the joint dispersal of mycorrhizal fungi and plant diaspores by
 954 birds. *New Phytologist* **222**: 1054–1060.
- 955 * CSATA, E., CZEKES, Z., EROS, K., NÉMET, E., HUGHES, M., CSOSZ, S. & MARKÓ, B. (2013).

- 956 Comprehensive survey of Romanian myrmecoparasitic fungi: New species, biology and
957 distribution. *North-Western Journal of Zoology* **9**, 23–29.
- 958 * CSÖSZ, S., LAPEVA-GJONOVA, A. & MARKÓ, B. (2012). New data on the geographical
959 distribution and host utilization of the entomopathogenic fungus *Myrmicinosporidium*
960 *durum*. *Journal of Insect Science* **12**, 1–5.
- 961 * CUESTA-SEGURA, A.D. & GARCIA GARCIA, F. (2016). Primera cita ibérica del mirmecófilo
962 *Myrmetes paykulli* Kanaar, 1979 (Coleoptera : Histeridae) con un nuevo hospedador para
963 la especie. *Boletín de la Sociedad Entomológica Aragonesa* **59**, 295–296.
- 964 * CUESTA-SEGURA, A.D., GARCÍA GARCÍA, F., CATARINEU, C., GARCIA-TEJERO, S. &
965 ESPADALER, X. (2018). Actualización de la distribución y hospedadores de la hormiga
966 parásita social *Teleutomyrmex schneideri* Kutter, 1959 en la Península Ibérica
967 (Hymenoptera : Formicidae). *Boletín de la Sociedad Entomológica Aragonesa (S.E.A.)* **63**,
968 225–239.
- 969 * CUSHING, P.E. (1997). Myrmecomorphy and myrmecophily in spiders: a review. *Florida*
970 *Entomologist* **80**, 165–193.
- 971 * CUSHING, P.E. (2012). Spider-ant Associations : an updated review of myrmecomorphy,
972 myrmecophily, and myrmecophagy in spiders. *Psyche: A Journal Of Entomology* **2012**,
973 151989.
- 974 * CZECHOWSKI, W. (1976). Competition between *Formica exsecta* Nyl. and *Formica*
975 *pressilabris* Nyl. (Hymenoptera, Formicidae). *Annales Zoologici* **33**, 273–286.
- 976 DARWIN, C. (1859). *On the Origin of the Species, or the Preservation of Favoured Races in the*
977 *Struggle for Life*. John Murray, London, UK.
- 978 DÁTILLO, W., GUIMARÃES JR., P.R. & IZZO, T.J. (2013). Spatial structure of ant-plant

979 mutualistic networks. *Oikos* **122**, 1643–1648.

980 * DE FERRER ANDREU, J., SÁEZ BOLAÑO, J.A. & BLANCO VILLERO, J.M. (2008). Los Histeridae
 981 (Coleoptera) de la sierra de Tudía (Badajoz, Extremadura, España). *Boletín de la Sociedad*
 982 *entomológica Aragonesa. S.E.A.* **43**, 409–415.

983 * DEKONINCK, W., LOCK, K. & JANSSENS, F. (2007). Acceptance of two native myrmecophilous
 984 species, *Platyarthrus hoffmannseggii* (Isopoda : Oniscidea) and *Cyphoderus albinus*
 985 (Collembola : Cyphoderidae) by the introduced invasive garden ant *Lasius neglectus*
 986 (Hymenoptera : Formicidae) in Belgium. *European Journal of Entomology*. **104**, 159–161.

987 DE LAENDER, F., ROHR, J.R., ASHAUER, R., BAIRD, D.J., BERGER, U., EISENHAUER, N., GRIMM,
 988 V., HOMMEN, U., MALTBY, L., MELIÀN, C.J. & POMATI, F., ROESSINK, I., RADCHUK, V. &
 989 VAN DEN BRINK, P.J (2016). Reintroducing environmental change drivers in
 990 biodiversity–ecosystem functioning research. *Trends in ecology & evolution* **31**, 905–
 991 915.

992 DEMARTIN, H. (2018). Hidden diversity of brain worms in zombie ants. The Pennsylvania State
 993 University Schreyer Honors College.

994 * DE MEIJERE, J.C.H. (1940) Eine neue myrmecophile *Phyllomyza*, *Phyllomyza pallida* n. sp.
 995 *Entomologische Berichten (Amsterdam)* **10**, 222–223.

996 * DEPA, Ł. (2010). Morphological variability within *Anoecia furcata* Theobald 1915
 997 /Hemiptera, Aphididae, Anoeciinae/. *Aphids and other Hemipterous Insects* **16**, 13–19.

998 * DEPA, Ł. (2012). Abundance of *Stomaphis graffii* Cholod. (Hemiptera) on maple trees in
 999 Poland. *Central European Journal of Biology*. **7**, 284–287.

1000 * DEPA, Ł., KANTURSKI, M., JUNKIERT, Ł. & WIECZOREK, K. (2015). Giant females vs dwarfish
 1001 males of the genus *Stomaphis* Walker (Hemiptera : Aphididae) – an aphid example of the
 1002 ongoing course to permanent parthenogenesis ? *Arthropod Systematics & Phylogeny* **73**,

- 1003 19–40.
- 1004 * DEPA, Ł., MRÓZ, E. & BROZEK, J. (2013). Description of the oviparous female and new
 1005 information on the biology of the rare aphid *Stomaphis radicola* Hille Ris Lambers 1947
 1006 (Hemiptera, Aphidoidea). *Entomologica Fennica* **24**, 100–106.
- 1007 * DEPA, Ł., MRÓZ, E., BUGAJ-NAWROCKA, A. & ORCZEWSKA, A. (2016). Do ants drive
 1008 speciation in aphids? A possible case of ant-driven speciation in the aphid genus *Stomaphis*
 1009 Walker (Aphidoidea, Lachninae). *Zoological Journal of the Linnean Society* **179**, 41–61.
- 1010 * DEPA, Ł., MRÓZ, E. & SZAWARYN, K. (2012). Molecular identity of *Stomaphis quercus*
 1011 (Hemiptera: Aphidoidea: Lachnidae) and description of a new species. *European Journal*
 1012 *of Entomology* **109**, 435–444.
- 1013 * DEPA, Ł. & WOJCIECHOWSKI, W. (2008). Ant-root aphid relations in different plant
 1014 associations. *Polish Journal of Entomology* **77**, 151–163.
- 1015 * DISNEY, R.H.L. (2000). Revision of European *Pseudaceton* Coquillett (Diptera, Phoridae).
 1016 *Bonner zoologische Beiträge*. **49**, 79–91.
- 1017 DONISTHORPE, H.S.J.K. (1927). *The guests of British ants, their habits and life-histories*.
 1018 George Routledge and Sons, London.
- 1019 * DOUWES, P., JESSEN, K. & BUSCHINGER, A. (1988). *Epimyrma alderzi* sp. n. (Hymenoptera:
 1020 Formicidae) from Greece: morphology and life history. *Entomologica scandinavica* **19**,
 1021 239–249.
- 1022 * DRANSFIELD, B., PHIL, D. & BRIGTHWELL, B. (2017). Aphids.
 1023 http://influentialpoints.com/Gallery/Aphid_genera.htm.
- 1024 * DRANSFIELD, R.D. & BRIGHTWELL, R. (2015). Aphids at Dundreggan, Scotland: July 2015
 1025 survey, 93.

- 1026 * DUPONT, S. & PAPE, T. (2007). Fore tarsus attachment device of the male scuttle fly,
1027 *Aenigmatias lubbockii*. *Journal of insect science* **7**, 1–8.
- 1028 * DURAK, R., WĘGRZYN, E. & LENIOWSKI, K. (2016). When a little means a lot-slight daily
1029 cleaning is crucial for obligatory ant-tended aphids. *Ethology Ecology and Evolution* **28**,
1030 20–29.
- 1031 * D' URSO, V. & MIFSUD, D. (2012). A preliminary account of the Auchenorrhyncha of the
1032 Maltese Islands (Hemiptera). *Bulletin of the Entomological Society of Malta* **5**, 57-72.
- 1033 * EASTOP, V.F. (1953). A study of the Tramini (Homoptera-Aphididae). *Transactions of the*
1034 *Royal Entomological Society of London* **104**, 385–415.
- 1035 EDWARDS, D.D. & MALCOLM, M. F. (2006). Host specificity among *Unionicola* spp. (Acari :
1036 Unionicolidae) parasitizing freshwater mussels. *The Journal of Parasitology* **92**, 977–983.
- 1037 ELIZALDE, L., PATROCK, R.J.W., DISNEY, R.H.L. & FOLGARAIT, P.J. (2018). Spatial and
1038 temporal variation in host–parasitoid interactions: leafcutter ant hosts and their phorid
1039 parasitoids. *Ecological Entomology* **43**, 114–125.
- 1040 ELMES, G.W., BARR, B., THOMAS, J.A. & CLARKE, R.T. (1999). Extreme host specificity by
1041 *Microdon mutabilis* (Diptera : Syrphidae), a social parasite of ants. *Proceedings of the*
1042 *Royal Society London B* **266**, 447–453.
- 1043 * ELTON, B.Y.C. (1932). Territory among wood ants (*Formica rufa* L.) at Picket Hill. *Journal*
1044 *of Animal Ecology* **1**, 69–76.
- 1045 * ESPADALER, X. & OLMO-VIDAL, J.M. (2011). The myrmecophilic cricket *Myrmecophilus* in
1046 Spain (Orthoptera, Myrmecophilidae). *Sociobiology* **57**, 321–328.
- 1047 ESPADALER, X. & SANTAMARIA, S. (2012). Ecto- and endoparasitic fungi on ants from the
1048 Holarctic Region. *Psyche: A Journal of Entomology* **2012**, 168478.

- 1049 EVANS, G.O. & TILL, W.M. (1966). Studies on the British Dermanyssidae (Acari:
1050 Mesostigmata) Part II. Classification. *Bulletin of The British Museum (Natural History)*
1051 *Zoology* **14**, 107–370.
- 1052 EZENWA, V.O., PRICE, S.A., ALTIZER, S., VITONE, N.D. & COOK, K.C. (2006). Host traits and
1053 parasite species richness in even and odd-toed hoofed mammals, Artiodactyla and
1054 Perissodactyla. *Oikos* **115**, 526–536.
- 1055 * FAIN, A. (1987). Observations on the hypopi of the genus *Forcellinia* Oudemans, 1924 (Acari,
1056 Acaridae). *Bulletin S.R.B.E/K.B.V.E.* **57**, 111–120.
- 1057 * FAIN, A. & CHMIELEWSKI, W. (1987). The phoretic hypopi of two acarid mites described from
1058 ants' nest: *Tyrophagus formicetorum* Volgin, 1948 and *Lasioacarus nidicolus* Kadzhaja
1059 and Sevastianov, 1967. *Acarologia* **28**, 53–61.
- 1060 FELIU, C., RENAUD, F., CATZEFLIS, F., HUGOT, J.-P., DURAND, P. & MORAND, S. (1997). A
1061 comparative analysis of parasite species richness of Iberian rodents. *Parasitology* **115**,
1062 453–466.
- 1063 * FIEDLER, K. (2006). Ant-associates of Palaearctic lycaenid butterfly larvae (Hymenoptera:
1064 Formicidae; Lepidoptera: Lycaenidae) - a review. *Myrmecological News.* **9**, 77–87.
- 1065 FONTAINE, C., GUIMARÃES JR., P.R., KÉFI, S., LOEUILLE, N., MEMMOTT, J., VAN DER PUTTEN,
1066 W.H., VAN VEEN, F.J.F. & THÉBAULT, E. (2011). The ecological and evolutionary
1067 implications of merging different types of networks. *Ecology Letters* **14**, 1170–1181.
- 1068 * FRANC, V. (1992). Myrmecophilous beetles of Slovakia with special reference to their
1069 endangerment and perspectives for protection. *Acta Universitatis Carolinae, Biologica* **36**,
1070 299–324.
- 1071 * FRANC, V. (2005). Prevailing trophic relations between spiders (Araneae) and ants
1072 (Formicoidea) in Slovakia. Contributions to Soil Zoology in Central Europe II. Tajovský,

- 1073 K., Schlaghamerský, J. & Pižl, V. (eds.): 41-45
- 1074 * FRANC, V., MAJZLAN, O., KRIŠTÍN, A. & WIEZIK, M. (2015). On the distribution and ecology
 1075 of the ant cricket (*Myrmecophilus acervorum*) (Orthoptera: Myrmecophilidae) in
 1076 Slovakia. *Proceedings of the conference «Roubal's Days I», Banská Bystrica, 27. 1. 2015,*
 1077 40–50.
- 1078 FRECKLETON, R.P., HARVEY, P.H. & PAGEL, M. (2002). Phylogenetic analysis and
 1079 comparative data: a test and review of evidence. *The American Naturalist* **160**, 712–726.
- 1080 * FREUDE, H., HARDE, K.W. & LOHSE, G.A. (1964). *Käfer Mitteleuropas Bd 4. Staphylinidae I*
 1081 (*Micropeplinae bis Tachyporinae*). Krefeld: Goecke et Evers Verlag.
- 1082 * FREUDE, H., HARDE, K.W. & LOHSE, G.A. (1974). *Käfer Mitteleuropas. Bd 5. Staphylinidae*
 1083 *II (Hypocypharinae und Aleocharinae), Pselaphidae*. Krefeld: Goecke et Evers Verlag.
- 1084 * GARCIA GARCIA, F. (2013). *Lasius myops*, Forel, 1894, nuevo hospedador del grillo
 1085 mirmecófilo *Myrmecophilus acervorum* (Panzer, 1799) (Orthoptera : Myrmecophilidae).
 1086 *Iberomyrmex* **5**, 9–10.
- 1087 * GARCÍA GARCÍA, F. & CUESTA-SEGURA, A.D. (2018). Coleópteros mirmecófilos ibéricos :
 1088 primeras citas y hospedadores (Coleoptera : Nitidulidae, Staphylinidae). *Boletín de la*
 1089 *Sociedad Entomológica Aragonesa* **63**, 307–309.
- 1090 * GARCÍA GARCÍA, F., CUESTA-SEGURA, A.D. & ESPADALER, X. (2017). Nuevas citas del
 1091 género *Chalepoxenus menozzi*, 1923 y actualización de su distribución y hospedadores
 1092 para la península Ibérica (Hymenoptera: Formicidae). *Iberomyrmex* **9**, 14–24.
- 1093 * GARCÍA GARCÍA, F. & ESPADALER, X. (2018). *Pandora formicae* (Humber y Balazi) Humber
 1094 1989 (Entomophthoromycota: Entomophthorales) en la Península Ibérica, con *Formica*
 1095 *pressilabris* Nylander, 1846 (Hymenoptera: Formicidae) como nuevo hospedante.

- 1096 *Iberomyrmex* **10**, 13–18.
- 1097 * GIEHR, J., HEINZE, J. & SCHREMPF, A. (2015). The ant *Cardiocondyla elegans* as host of the
 1098 enigmatic endoparasitic fungus *Myrmecinosporidium durum*. *Psyche: A Journal of*
 1099 *Entomology* **2015**, 364967
- 1100 * GJONOV, I. (2002). Data on the bionomics and distribution of *Balcanocerus balcanicus*
 1101 (Horváth, 1903). (Homoptera, Cicadina, Cicadellidae) in Bulgaria. *Entomological Review*
 1102 **82**, 641-642.
- 1103 * GJONOV, I. & LAPEVA-GJONOVA, A. (2013). New data on ant-attendance in leafhoppers
 1104 (Hemiptera: Cicadellidae). *North-Western Journal of Zoology* **9**, 433–437.
- 1105 GLASIER, J.R.N., POORE, A.G.B. & ELDRIDGE, D.J. (2018). Do mutualistic associations have
 1106 broader host ranges than neutral or antagonistic associations? A test using
 1107 myrmecophiles as model organisms. *Insectes Sociaux* **65**, 639–648.
- 1108 * GÓMEZ DURÁN, J.-M. (2012). Moscas fóridas en un árbol caído. *Microselia southwoodi*
 1109 Disney (Diptera, Phoridae): primera cita para España y notas sobre su comportamiento
 1110 ovipositor.
- 1111 * GÓMEZ, K. & ESPADALER, X. (2008). Dos especies del género *Lasius* Fabricius 1804
 1112 (Hymenoptera, Formicidae) nuevas para la Península Ibérica. *Orsis* **23**, 163–166.
- 1113 * GÓMEZ, K., ESPADALER, X. & SANTAMARIA, S. (2016). Ant-fungus interactions: *Laboulbenia*
 1114 *camponoti* Batra in Italy and a new host for *L. formicarum* Thaxter (Fungi: Ascomycota,
 1115 laboulbeniales). *Sociobiology* **63**, 950–955.
- 1116 * GONÇALVES, C., PATANITA, I. & ESPADALER, X. (2012). Substantial, and significant,
 1117 expansion of ant hosts range for *Myrmecinosporidium Hölldobler*, 1933 (Fungi). *Insectes*
 1118 *Sociaux* **59**, 395–399.

- 1119 * GÖSSWALD, K. (1950) Pflege des Ameisenparasiten *Tamiclea globula* Meig. (Dipt.) durch
 1120 den Wirt mit Bemerkungen über den Stoffwechsel in der parasitierten Ameise. In
 1121 *Verhandlungen der Deutschen Zoologen (Mainz, 1949)* pp.256-264.
- 1122 GRILLI, J., BARABÁS, G., MICHALSKA-SMITH, M. J., & ALLESINA, S. (2017). Higher-order
 1123 interactions stabilize dynamics in competitive network models. *Nature* **548**, 210–213.
- 1124 * GROSSO-SILVA, J. & SERRANO, A.R.M. (2000). Registos interessantes de Coleopteros
 1125 (Insecta, Coleoptera) para Portugal (2ª nota). *Boletín de la S.E.A.* **27**, 1–5.
- 1126 GUEGAN, J.-F, LAMBERT, A., LEVEQUE, C., COMBES, C. & EUZET, L. (1992). Can host body size
 1127 explain the parasite species richness in tropical freshwater fishes? *Oecologia* **90**, 197–204.
- 1128 GUIMARÃES JR., P. R., JORDANO, P. & THOMPSON, J. N. (2011). Evolution and coevolution in
 1129 mutualistic networks. *Ecology Letters*, **14**, 877–885.
- 1130 GUIMARÃES JR., P.R., RICO-GRAY, V., DOS REIS, S.F. & THOMPSON, J.N. (2006). Asymmetries
 1131 in specialization in ant-plant mutualistic networks. *Proceedings of the Royal Society B:*
 1132 *Biological Sciences* **273**, 2041–2047.
- 1133 HAIRSTON, N.G., SMITH, F.E. & SLOBODKIN, L.B. (1960). Community structure, population
 1134 control, and competition. *The American Naturalist* **94**, 421–425.
- 1135 * HANSEN, L.O. & ROLSETH, K. (2004). The black willow aphid *Pterocomma salicis* (L .,
 1136 1758). (Hemiptera, Aphididae) in Norway. *The Norwegian Journal of Entomologia* **51**,
 1137 189–191.
- 1138 * HARRISON, T. (2016). *Dinarda dentata* (Gravenhorst, 1806). (Coleoptera : Staphylinidae :
 1139 Aleocharinae) new to Bulgaria and an additional site for *Lomechusoides strumosus*
 1140 (Fabricius, 1793). in Bulgaria. *ZooNotes* **86**, 1–4.
- 1141 * HÁVA, J. (2015). *World Catalogue of Insects. Volume 13. Dermestidae (Coleoptera)*. Brill
 1142 Academic Pub.

- 1143 * HEIE, O.E. (1980). The Aphidoidea (Hemiptera) of Fennoscandia and Denmark. I General
1144 Part. The Families Mindaridae, Hormaphididae, Thelaxidae, Anoeciidae, and
1145 Pemphigidae. Fauna Entomologica Scandinavica, Scandinavian Science Press, Vinderup,
1146 Denmark.
- 1147 HIMLER, A.G., ADACHI-HAGIMORI, T., BERGEN, J.E., KOZUCH, A., KELLY, S.E., TABASHNIK,
1148 B.E., CHIEL, E., DUCKWORTH, V.E., DENNEHY, T.J., ZCHORI-FEIN, E. & HUNTER, M.S.
1149 (2011). Rapid spread of a bacterial symbiont in an invasive whitefly is driven by fitness
1150 benefits and female bias. *Science* **332**, 254–256.
- 1151 * HLAVÁČ, P. (2005). Revision of the myrmecophilous genus *Lomechusa* (Coleoptera:
1152 Staphylinidae: Aleocharinae). *Sociobiology* **46**, 203–250.
- 1153 * HLAVÁČ, P., BEKCHIEV, R., RŮŽIČKA, J. & LACKNER, T. (2007). Contribution to the
1154 knowledge of myrmecophilous beetles (Coleoptera) of Bulgaria. *Acta Societatis*
1155 *Zoologicae Bohemicae* **71**, 127–132.
- 1156 * HLAVÁČ, P., NEWTON, A.F. & MARUYAMA, M. (2011). World catalogue of the species of the
1157 tribe Lomechusini (Staphylinidae: Aleocharinae). *Zootaxa* **151**, 1–151.
- 1158 HÖLLDOBLER, B. & WILSON, E.O. (1990). *The ants*. Harvard University Press, Cambridge,
1159 Massachusetts.
- 1160 * HORNUNG, E., VILISICS, F. & TARTALLY, A. (2005). Occurrence of *Platyarthrus schoblii*
1161 (Isopoda, Oniscidea) and its ant hosts in Hungary. *European Journal of Soil Biology* **41**,
1162 129–133.
- 1163 * HUGHES, T.E. (1959). *Mites or the Acari*. The Athlone Press, University of London, London.
- 1164 HUGHES, D.P., PIERCE, N.E. & BOOMSMA, J.J. (2008). Social insect symbionts: evolution in
1165 homeostatic fortresses. *Trends in Ecology and Evolution* **23**, 672–677.

1166 INGS, T.C., MONTOYA, J.M., BASCOMPTE, J., BLÜTHGEN, N., BROWN, L., DORMANN, C.F.,
 1167 EDWARDS, F., FIGUEROA, D., JACOB, U., JONES, J.I., LAURIDSEN, R.B., LEDGER, M.E.,
 1168 LEWIS, H.M., OLESEN, J.M., VAN VEEN, F.J.F., *ET AL.* (2009). Ecological networks –
 1169 beyond food webs. *Journal of Animal Ecology* **78**, 253–269.

1170 IVENS, A.B.F., von BEEREN, C., BLÜTHGEN, N. & KRONAUER, D.J.C. (2016). Studying the
 1171 complex communities of ants and their symbionts using ecological network analysis.
 1172 *Annual Review of Entomology* **61**, 353–371.

1173 IVENS, A.B.F., GADAU, A., KIERS, E.T. & KRONAUER, D.J.C. (2018). Can social partnerships
 1174 influence the microbiome? Insights from ant farmers and their trophobiont mutualists.
 1175 *Molecular Ecology* **27**, 1898–1914.

1176 * IUCN (2017). IUCN Red List of Threatened Species. *Version 2017.3*. www.iucnredlist.org.

1177 JACQUEMYN, H., MERCKX, V., BRYN, R., TYTECA, D., CAMMUE, B.P.A., HONNAY, O. &
 1178 LIEVENS, B. (2011). Analysis of network architecture reveals phylogenetic constraints on
 1179 mycorrhizal specificity in the genus *Orchis* (Orchidaceae). *New Phytologist* **192**, 518–528.

1180 JAENIKE, J., UNCKLESS, R., COCKBURN, S.N., BOELIO, L.M. & PERLMAN, S.J. (2010).
 1181 Adaptation via symbiosis: recent spread of a drosophila defensive symbiont. *Science*
 1182 **329**, 212–215.

1183 JANET, C. (1897). *Études sur les fourmis, les guêpes et les abeilles. Note 14: Rapports des*
 1184 *animaux myrmécophiles avec les fourmis*. Ducourtieux, Limoges.

1185 * JANSEN, G., SAVOLAINEN, R. & VEPSÄLÄINEN, K. (2010). Molecular phylogenetics and
 1186 evolution phylogeny, divergence-time estimation, biogeography and social parasite–host
 1187 relationships of the Holarctic ant genus *Myrmica* (Hymenoptera : Formicidae). *Molecular*
 1188 *Phylogenetics and Evolution* **56**, 294–304.

1189 * JOHANSSON, T. & GIBB, H. (2012). Forestry alters foraging efficiency and crop contents of
 1190 aphid-tending red wood ants, *Formica aquilonia*. *PLoS One* **7**, e32817

- 1191 JOHNSON, C. A., & BRONSTEIN, J. L. (2019). Coexistence and competitive exclusion in
1192 mutualism. *Ecology* **100**, e02708.
- 1193 * JUNKER, E.A. (1997). Untersuchungen zur Lebensweise und Entwicklung von *Myrmecophilus*
1194 *acervorum* (PANZER, 1799). (Saltatoria, Myrmecophilidae). *Articulata* **12**, 93–106.
- 1195 * JUNKER, E.A. & RATSCHKER, U.M. (2000). Zur Verbreitung der Ameisengrille,
1196 *Myrmecophilus acervorum* (Panzer, 1799)., in Sachsen (Insecta; Ensifera;
1197 Myrmecophilidae). *Faunistische Abhandlungen Staatliches Museum für Tierkunde*
1198 *Dresden* **22**, 11–21.
- 1199 KAMIYA, T., O'DWYER, K., NAKAGAWA, S. & POULIN, R. (2014). What determines species
1200 richness of parasitic organisms ? A meta-analysis across animal, plant and fungal hosts.
1201 *Biological Reviews* **89**, 123–134.
- 1202 * KANTURSKI, M., KARCZ, J., KASZYCA, N. & ŁUKASZ, D. (2017). Perianal structures in
1203 myrmecophilous subterranean aphids (Insecta: Hemiptera: Aphididae)–Comparative
1204 morphology of trophobiotic organ with its first description in Lachninae. *Arthropod*
1205 *Structure and Development* **46**, 496–507.
- 1206 * KARSA, D., HALTRICH, A. & SZATHMÁRY, E. (2004). Two new species in the Hungarian fauna:
1207 *Stomaphis mordvilko* Hille Ris Lambers and *Stomaphis juglandis* Petrovič (Lachnidae,
1208 Homoptera). *Acta Phytopathologica et Entomologica Hungarica* **39**, 405–407.
- 1209 KASPARI, M. & VARGO, E.L. (1995). Colony size as a buffer against seasonality: Bergmann's
1210 rule in social insects. *The American Naturalist* **145**, 610–632.
- 1211 KAWAKITA, A., OKAMOTO, T., GOTO, R. & KATO, M. (2010). Mutualism favours higher host
1212 specificity than does antagonism in plant-herbivore interaction. *Proceedings of the Royal*
1213 *Society B: Biological Sciences* **277**, 2765–2774.
- 1214 * KHAUSTOV, A.A. (2014). A review of myrmecophilous mites of the family microdispidae

- 1215 (Acari, Heterostigmatina) of Western Siberia. *ZooKeys* **28**, 13–28.
- 1216 * KHAUSTOV, A.A. (2015). Myrmecophilous pygmephoroid mites (Acari: Pygmephoroidae)
- 1217 associated with *Lasius flavus* (Hymenoptera: Formicidae) in Russia. *Zootaxa* **4044**, 345–
- 1218 370.
- 1219 * KHAUSTOV, A.A. (2016). Myrmecophilous pygmephoroid mites (Acari: Pygmephoroidae)
- 1220 associated with *Lasius fuliginosus* (Hymenoptera: Formicidae) in Western Siberia, Russia.
- 1221 *International Journal of Acarology* **42**, 92–105.
- 1222 * KHAUSTOV, A.A. & TOLSTIKOV, A. V (2016). The diversity, mite communities, and host
- 1223 specificity of pygmephoroid mites (Acari: Pygmephoroidae) associated with ants in
- 1224 western Siberia, Russia. *Acarina* **24**, 113–136.
- 1225 * KHAUSTOV, A.A. & TRACH, V.A. (2013). New and little-known species of myrmecophilous
- 1226 mites of the genus *Petalomium* (Acari: Heterostigmata: Neopygmephoridae) from
- 1227 Ukraine. *Acarina* **21**, 43–52.
- 1228 * KIRAN, K., KARAMAN, C., LAPEVA-GJONOVA, A. & AKSOY, V. (2017). Two new species of
- 1229 the ‘ultimate’ parasitic ant genus *Teleutomyrmex* Kutter, 1950 (Hymenoptera: Formicidae)
- 1230 from the Western Palaearctic. *Myrmecological News* **25**, 145–155.
- 1231 KISTNER, D.H. (1979). Social and evolutionary significance of social insect symbionts. In
- 1232 *Social Insects, Volume I* (ed H.R. HERMAN), pp. 339–413. Academic Press.
- 1233 KISTNER, D.H. (1982). The social insects’ bestiary. In *Social Insects, Vol III* (ed H.R.
- 1234 HERMAN), pp. 1–244. Academic Press.
- 1235 * KLIMES, P., BOROVANSKA, M., PLOWMAN, N.S. & LEPONCE, M. (2018). How common is
- 1236 trophobiosis with hoppers (Hemiptera: Auchenorrhyncha) inside ant nests (Hymenoptera:
- 1237 Formicidae)? Novel interactions from New Guinea and a worldwide overview.
- 1238 *Myrmecological News* **2018**, 31–45.

- 1239 * KÖHLER, A. (2010). Notes on the biology of *Holoplagia lucifuga* (Loew) (Scatopsidae,
1240 Diptera). *Studia dipterologica* **17**, 81–90.
- 1241 * KORGE, H. (2005). Rote Liste und Gesamtartenliste der Kurzflügelkäfer (Coleoptera :
1242 Staphylinidae) von Berlin. In: Der Landesbeauftragte für Naturschutz und
1243 Landschaftspflege/Senatsverwaltung für Stadtentwicklung (Hrsg.). Rote Listen der
1244 gefährdeten Pflanzen und Tiere von Berlin, 1-11.
- 1245 * KOSZTARAB, M. & KOZÁR, F. (1988). *Scale Insects of Central Europe*. Akadémiai Kiadó,
1246 Budapest.
- 1247 KRONAUER, D.J.C. & PIERCE, N.E. (2011). Myrmecophiles. *Current Biology* **21**, 208–209.
- 1248 * KUNKEL, H. (1973). Die Kotabgabe der Aphiden (Aphidina, Hemiptera) unter Einfluß von
1249 Ameisen. *Bonner Zoologische Beiträge* **24**, 105–121.
- 1250 KURIS, A.M., BLAUSTEIN, A.R. & ALIÓ, J.J. (1980). Hosts as islands. *The American Naturalist*
1251 **116**, 570–586.
- 1252 * LACHAUD, J.-P. & PÉREZ-LACHAUD, G. (2012). Diversity of species and behavior of
1253 Hymenopteran parasitoids of ants: a review. *Psyche: A Journal of Entomology* **2012**,
1254 134746.
- 1255 * LAPEVA-GJONOVA, A. (2004). Pselaphinae (Coleoptera: Staphylinidae) from ant nests
1256 (Hymenoptera: Formicidae) in Southwestern Bulgaria. *Acta Zoologica Bulgarica* **56**, 69–
1257 73.
- 1258 * LAPEVA-GJONOVA, A. (2013). Ant-Associated Beetle Fauna in Bulgaria : A Review and new
1259 data. *Psyche: A Journal of Entomology* **2013**, 242037.
- 1260 * LAPEVA-GJONOVA, A. (2014). *Cataglyphis aenescens* - A newly discovered ant host of the
1261 fungal parasite *Myrmecinosporidium durum*. *Bulgarian Journal of Agricultural Science*

- 1262 **20**, 157–159.
- 1263 * LAPEVA-GJONOVA, A., KIRAN, K. & AKSOY, V. (2012). Unusual ant hosts of the socially
 1264 parasitic ant *Anergates atratulus* (Schenck, 1852). (Hymenoptera, Formicidae). *Psyche: A*
 1265 *Journal of Entomology* **2012**, 391525.
- 1266 * LAPEVA-GJONOVA, A. & LIEFF, O. (2012). Ant-associated rove beetles (Coleoptera:
 1267 Staphylinidae) in Bulgaria. *Acta Entomologica Slovenica* **20**, 73–84.
- 1268 * LAPEVA-GJONOVA, A. & RÜCKER, W.H. (2011). Latridiidae and Endomychidae beetles
 1269 (Coleoptera) from ant nests in Bulgaria. *Latridiidae* **8**, 5–8.
- 1270 * LARSSON, S. (1943). *Danmarks Fauna: Bd. 49 Myrer*. I Kommission hos G.E.C. Gads Forlag
 1271 Kobenhavn.
- 1272 * LEBAS, C., GALKOWSKI, C., WEGNEZ, P., ESPADALER, X. & BLATRIX, R. (2015). La diversité
 1273 exceptionnelle des fourmis du Mont Coronat (Pyrénées Orientales), et *Temnothorax*
 1274 *gredosi* (Hymenoptera, Formicidae) nouveau pour la France. *R.A.R.E.* **24**, 24–33.
- 1275 * LEHOUCK, V., BONTE, D., DEKONINCK, W. & MAELFAIT, J.-P. (2004). Trophobiotic
 1276 relationships between ants (Hymenoptera: Formicidae) and Tettigometridae (Hemiptera:
 1277 Fulgoromorpha) in the grey dunes of Belgium. *European Journal of Entomology* **101**, 547-
 1278 553.
- 1279 LEIBOLD, M.A., HOLYOAK, M., MOUQUET, N., AMARASEKARE, P., CHASE, J.M., HOOPES,
 1280 M.F., HOLT, R.D., SHURIN, J.B., LAW, R., TILMAN, D., LOREAU, M. & GONZALEZ, A.
 1281 (2004). The metacommunity concept: a framework for multi-scale community ecology.
 1282 *Ecology Letters* **7**, 601–613.
- 1283 * LE MASNE, G. & TOROSSIAN, C. (1965). Observations sur le comportement du coléoptère
 1284 myrmécophile *Amorphocephalus coronatus* Germar (Brenthidae) hôte des *Camponotus*.

- 1285 *Insectes Sociaux* **12**, 185–194.
- 1286 * LESNE P. (1905). Les relations des fourmis avec les hémiptères homoptères de la famille des
 1287 fulgorides; domestication des Tettigometra. *Bulletin de la Société Entomologique de*
 1288 *France* **10**, 162–164.
- 1289 * LENCINA, J.L., TORRES, J.L., BAENA, M., ANDÚJAR, C., GALLEG0, D., GONZÁLEZ, E. &
 1290 ZUZARTE, A.J. (2011). Notas sobre *Amphotis* Erichson, 1843 Ibéricos (Coleoptera:
 1291 Nitidulidae). *Boletín de la Sociedad Entomológica Aragonesa (S.E.A.)* **49**, 149–152.
- 1292 * LENOIR, A., CHALON, Q., CARVAJAL, A., RUEL, C., BARROSO, Á., LACKNER, T. & BOULAY,
 1293 R. (2012). Chemical integration of myrmecophilous guests in *Aphaenogaster* ant nests.
 1294 *Psyche: A Journal of Entomology* **2012**, 840860.
- 1295 * LENOIR, A., HÁVA, J., HEFETZ, A., DAHBI, A., CERDÁ, X. & BOULAY, R. (2013). Chemical
 1296 integration of *Thorictus* myrmecophilous beetles into *Cataglyphis* ant nests. *Biochemical*
 1297 *Systematics and Ecology* **51**, 335–342.
- 1298 * LEPPÄNEN, J., SEPPÄ, P., VEPSÄLÄINEN, K. & SAVOLAINEN, R. (2016). Mating isolation
 1299 between the ant *Myrmica rubra* and its microgynous social parasite. *Insectes Sociaux* **63**,
 1300 79–86.
- 1301 LICHSTEIN, J.W. (2007). Multiple regression on distance matrices : a multivariate spatial
 1302 analysis tool. *Plant Ecology* **188**, 117–131.
- 1303 LINDENFORS, P., NUNN, C.L., JONES, K.E., CUNNINGHAM, A.A., SECHREST, W. & GITTLEMAN,
 1304 J.L. (2007). Parasite species richness in carnivores: effects of host body mass, latitude,
 1305 geographical range and population density. *Global Ecology and Biogeography* **16**, 496–
 1306 509.
- 1307 * LOMPE, A. (2017). die Käfer Europas. www.coleo-net.de [accessed 20 July 2018].

- 1308 LUIS, A.D., O'SHEA, T.J., HAYMAN, D.T.S., WOOD, J.L.N., CUNNINGHAM, A.A., GILBERT,
1309 A.T., MILLS, J.N. & WEBB, C.T. (2015). Network analysis of host-virus communities in
1310 bats and rodents reveals determinants of cross-species transmission. *Ecology Letters* **18**,
1311 1153–1162.
- 1312 * LUZE, G. (1901). Revision der europäischen und sibirischen Arten der Staphyliniden-
1313 Gattungen *Tachyporus* Grav. und *Lamprinus* Heer. *Verhandlungen der Kaiserlich-*
1314 *Königlichen Zoologisch-Botanischen Gesellschaft in Wien* **51**, 146–185.
- 1315 MACARTHUR, R.H. & WILSON, E.O. (1967). *The theory of island biogeography*. Princeton
1316 University Press.
- 1317 * MAHUNKA, S. (1967). Beiträge zur Kenntnis der Tschechoslowakischen Tarsonemini-Fauna.
1318 *Vestník Československé Společnosti Zoologické*. **31**, 240–244.
- 1319 * MAJZLAN, O. (2007). Faunistické příspěvky zo Slovenska (Coleoptera) 4. *Naturae Tutela* **12**,
1320 207–210.
- 1321 * MAŁOL, J., ŁAYDANOWICZ, J. & KŁOSIŃSKA, A. (2010). Myrmecophilous Parasitengona
1322 (Acari : Prostigmata) — Accident or Adaptation? *Annales Zoologici* **60**, 429–437.
- 1323 * MAŁAGOCKA, J., JENSEN, A.B. & EILENBERG, J. (2017). *Pandora formicae*, a specialist ant
1324 pathogenic fungus: New insights into biology and taxonomy. *Journal of Invertebrate*
1325 *Pathology* **143**, 108–114.
- 1326 * MALENOVSKÝ, I. & LAUTERER, P. (2012). Leafhoppers and planthoppers (Hemiptera:
1327 Auchenorrhyncha) of the Bílé Karpaty Protected Landscape Area and Biosphere Reserve
1328 (Czech Republic). *Acta Musei Moraviae, Scientiae biologicae* **96**, 155–322.
- 1329 * MARTINEZ GARCIA, A. (2011). *Amorphocephala coronata* (Germar 1817). (Coleoptera :
1330 Brentidae) en la provincia de Málaga, España. *Boletín de la SAE* **17**, 12–17.

- 1331 MARTINY, J.B.H., EISEN, J.A., PENN, K., ALLISON, S.D. & HORNER-DEVINE, M.C. (2011).
 1332 Drivers of bacterial β -diversity depend on spatial scale. *PNAS* **108**, 7850–7854.
- 1333 * MARUYAMA, M. (2006). Revision of the Palearctic species of the myrmecophilous genus
 1334 *Pella* (Coleoptera Staphylinidae, Aleocharinae). *National Science Museum Monographs*,
 1335 1–207.
- 1336 MASCHWITZ, U. & HÖLLDOBLER, B. (1970). Der Kartonnestbau bei *Lasius fuliginosus* Latr.
 1337 (Hym. Formicidae). *Zeitschrift für Vergleichende Physiologie* **66**, 176–189.
- 1338 * MATHES, I. & STROUHAL, H. (1954). Zur Ökologie und Biologie der Ameisenassel
 1339 *Platyarthrus hoffmannseggii*. *Zeitschrift für Morphologie und Ökologie der Tiere* **43**, 82–
 1340 93.
- 1341 * MAYORAL, J.G. & BARRANCO, P. (2010). The little known genus *Forania* (Acari, Prostigmata,
 1342 Erythraeidae). *Zootaxa* **2408**, 59–68.
- 1343 * MICHEL, E. (1942). Beiträge zur Kenntnis von *Lachnus* (*Pterochlorus*) *roboris* L., einer
 1344 wichtigen Honigtauerzeugerin an der Eiche. *Zeitschrift für Angewandte Entomologie* **29**,
 1345 243–281.
- 1346 * MIÑARRO, M., FERNÁNDEZ-MATA, G. & MEDINA, P. (2010). Role of ants in structuring the
 1347 aphid community on apple. *Ecological Entomology* **35**, 206–215.
- 1348 MOLERO-BALTANÁS, R., BACH DE ROCA, C., TINAUT, A., PÉREZ, J.D. & GAJU-RICART, M.
 1349 (2017). Symbiotic relationships between silverfish (Zygentoma: Lepismatidae,
 1350 Nicoletiidae) and ants (Hymenoptera: Formicidae) in the Western Palaearctic. A
 1351 quantitative analysis of data from Spain. *Myrmecological News* **24**, 107–122.
- 1352 * MONTESANTO, G., CARUSO, D. & LOMBARDO, B.M. (2008). Genetic variability in
 1353 parthenogenetic and amphigonic populations of *Platyarthrus aiasensis* Legrand from

- 1354 Sicily (Isopoda, Oniscidea). *Proceedings of the International Symposium of Terrestrial*
 1355 *Isopod Biology* **7**, 59–67.
- 1356 MOUGI, A. & KONDOH, M. (2012). Diversity of interaction types and ecological community
 1357 stability. *Science* **337**, 349–351.
- 1358 * MUIR, D.A. (1954). Ants, *Myrmica rubra* L. and *M. scabrinodis* Nylander as intermediate
 1359 hosts of a cestode. *Nature* **173**: 688-689.
- 1360 * MUIR, D.A. (1959). The ant-aphid-plant relationship in west Dumbartonshire. *Journal of*
 1361 *Animal Ecology* **28**, 133–140.
- 1362 * MULLER, C.B., ADRIAANSE, I.C.T., BELSHAW, R. & GODFRAY, H.C.J. (1999). The structure
 1363 of an aphid-parasitoid community. *Journal of Animal Ecology* **68**, 346–370.
- 1364 * NAPIERAŁA, A. & BŁOSZYK, J. (2013). Unstable microhabitats (merocenoses) as specific
 1365 habitats of *Uropodina* mites (Acari: Mesostigmata). *Experimental and Applied Acarology*
 1366 **60**, 163–180.
- 1367 NAVA, S. & GUGLIELMONE, A.A. (2013). A meta-analysis of host specificity in Neotropical
 1368 hard ticks (Acari: Ixodidae). *Bulletin of Entomological Research* **103**, 216–224.
- 1369 * NEMESTÓTHY, K.K. & MAHUNKA, S. (1981). Tarsoneminen (acari) aus dem
 1370 Naturschutzgebiet Wacholders „Borókás" bei Barcs, Ungarn. *Dunántúli Dolgozatok*
 1371 *Természettudományi soroz*a **2**, 157–166.
- 1372 NEWMAN, M.E.J. (2003). The structure and function of complex networks. *SIAM Review* **45**,
 1373 167–256.
- 1374 * NICKEL, H., HOLZINGER, W.E. & WACHMANN, E. (2008). Mitteleuropäische Lebensräume und
 1375 ihre Zikadenfauna (Hemiptera: Auchenorrhyncha). Universitätsbibliothek Johann
 1376 Christian Senckenberg.
- 1377 NOSIL, P. (2002). Transition rates between specialization and generalization in phytophagous

- 1378 insects. *Evolution* **56**, 1701–1706.
- 1379 * NOVGORODOVA, T.A. (2002). Study of adaptations of aphids (Homoptera, Aphidinea) to ants :
- 1380 Comparative analysis of myrmecophilous and nonmyrmecophilous species.
- 1381 *Entomological Review* **82**, 569–576.
- 1382 * NOVGORODOVA, T.A. (2003). Myrmecophilous complexes of aphids in forest and steppe areas
- 1383 of Novosibirsk Oblast'. *Euroasian Entomological Journal* **2**, 243–250.
- 1384 NOVGORODOVA, T.A. (2005). Red wood ants (Formicidae) impact on multi-species complexes
- 1385 of aphids (Aphididae) in the forest-park zone of Novosibirsk. *Euroasion Entomological*
- 1386 *Journal* **4**, 117–120.
- 1387 * NOVGORODOVA, T.A. (2015). Organization of honeydew collection by foragers of different
- 1388 species of ants (Hymenoptera: Formicidae): Effect of colony size and species specificity.
- 1389 *European Journal of Entomology* **112**, 688–697.
- 1390 * NOVGORODOVA, T.A. (2016). Peculiarities of trophobiotic interactions with aphids in slave-
- 1391 maker ant *Formica sanguinea* Latreille, 1798 (Hymenoptera, Formicidae). *Eurasian*
- 1392 *Entomological Journal* **15**, 108–114.
- 1393 * NOVGORODOVA, T.A. & GAVRILYUK, A. V. (2012). The degree of protection different ants
- 1394 (Hymenoptera: Formicidae) provide aphids (Hemiptera: Aphididae) against
- 1395 aphidophages. *European Journal of Entomology* **109**, 187–196.
- 1396 * NOVGORODOVA, T.A. & RYABININ, A.S. (2015). Trophobiotic associations between ants
- 1397 (Hymenoptera, Formicidae) and aphids (Hemiptera, Aphidomorpha) in South Zauralye.
- 1398 *Экология* **15**, 98–107.
- 1399 * NOVGORODOVA, T.A. & RYABININ, A.S. (2018). Ant–aphid relations in the south of Western
- 1400 Siberia (Hymenoptera: Formicidae; Hemiptera: Aphididae). *Arthropod-Plant Interactions*

- 1401 **12**, 369-376.
- 1402 NUNN, C.L., ALTIZER, S., JONES, K.E. & SECHREST, W. (2003). Comparative tests of parasite
1403 species richness in primates. *The American Naturalist* **162**, 597–614.
- 1404 * O’GRADY, A. & BREEN, J. (2011). Observations on mermithid parasitism (Nematoda:
1405 Mermithidae) in two species of *Lasius* ants (Hymenoptera: Formicidae). *Journal of*
1406 *Natural History* **45**, 2339–2345.
- 1407 * O’KEEFE, S. (2000). Ant-like stone beetles, ants, and their associations (Coleoptera:
1408 Scydmaenidae; Hymenoptera: Formicidae; Isoptera). *Journal of the New York*
1409 *Entomological Society* **108**, 273–303.
- 1410 OKUYAMA, T. & HOLLAND, J.N. (2008). Network structural properties mediate the stability of
1411 mutualistic communities. *Ecology Letters* **11**: 208–216.
- 1412 OLESEN, J.M., BASCOMPTE, J., DUPONT, Y.L. & JORDANO, P. (2007). The modularity of
1413 pollination networks. *Proceedings of the National Academy of Sciences of the United*
1414 *States of America* **104**, 19891–19896.
- 1415 * ÖZDEMİR, I., AKTAÇ, N., TOROS, S., KILINÇER, N. & GÜRKAN, O. (2008). Investigations of
1416 the associated between aphids and ants on wild plants in Ankara Province (Turkey). *Munis*
1417 *Entomology & Zoology Journal* **3**, 606–613.
- 1418 * PACLT, J. (1987). Ein Nachtrag zur geographischen Verbreitung von *Entomobrya*
1419 *myrmecophila* Reuter (Collembola). *Mitteilungen Zoologisches Museum Berlin* **63**, 317–
1420 319.
- 1421 PAGEL, M. (1999). Inferring the historical patterns of biological evolution. *Nature* **401**, 877–
1422 884.
- 1423 * PÄIVINEN, J., AHLROTH, P. & KAITALA, V. (2002). Ant-associated beetles of Fennoscandia
1424 and Denmark. *Entomologica Fennica* **13**, 20–40.

- 1425 PARACER, S. & AHMADJIAN, V. (2000). *Symbiosis: An introduction to biological associations*.
 1426 Oxford University Press, New York.
- 1427 PARMENTIER, T. (2019). Host following of an ant associate during nest relocation. *Insectes*
 1428 *Sociaux* **66**, 329–332.
- 1429 PARMENTIER, T. (2020). Guests of social insects. In *Encyclopedia of Social Insects* (ed C.K.
 1430 Starr). Springer, Cham. https://doi.org/10.1007/978-3-319-90306-4_164-1
- 1431 PARMENTIER, T., BOUILLON, S., DEKONINCK, W. & WENSELEERS, T. (2016a). Trophic
 1432 interactions in an ant nest microcosm: a combined experimental and stable isotope
 1433 ($\delta^{13}\text{C}/\delta^{15}\text{N}$) approach. *Oikos* **125**, 1182–1192.
- 1434 PARMENTIER, T., DEKONINCK, W., & WENSELEERS, T. (2016b). Survival of persecuted
 1435 myrmecophiles in laboratory nests of different ant species can explain patterns of host
 1436 use in the field (Hymenoptera: Formicidae). *Myrmecological News* **23**, 71–79.
- 1437 * PARMENTIER, T., DEKONINCK, W. & WENSELEERS, T. (2014) A highly diverse microcosm in
 1438 a hostile world: a review on the associates of red wood ants (*Formica rufa* group). *Insectes*
 1439 *Sociaux* **61**, 229–237.
- 1440 PARMENTIER, T., DEKONINCK, W. & WENSELEERS, T. (2017). Arthropods associate with their
 1441 red wood ant host without matching nestmate recognition cues. *Journal of Chemical*
 1442 *Ecology* **43**, 644–661.
- 1443 PARMENTIER, T., DE LAENDER, F., WENSELEERS, T. & BONTE, D. (2018). Prudent behavior
 1444 rather than chemical deception enables a parasite to exploit its ant host. *Behavioral*
 1445 *Ecology* **29**, 1225–1233.
- 1446 * PASSERA, L. (1974) Presence d'*Hexamermis* sp. (Nematoda, Mermithidae) dans les reines
 1447 vierges et les males de la fourmi *Pheidole pallidula* Nyl. (Formicidae, Myrmicinae).
 1448 *Bulletin de la Societe Zoologique de France* **99**, 315–324.
- 1449 * PECH, P. (2013). *Myrmica curvithorax* (Hymenoptera: Formicidae) in the Czech Republic: a

1450 contribution to the knowledge of its distribution and biology. *Klapalekiana* **49**, 197–204.

1451 PEDERSEN, A.B., ALTIZER, S., POSS, M., CUNNINGHAM, A.A. & NUNN, C.L. (2005). Patterns
1452 of host specificity and transmission among parasites of wild primates. *International*
1453 *Journal for Parasitology* **35**, 647–657.

1454 * PÉREZ-HIDALGO, N., ESPADALER, X. & NIETO, J.M. (2007). Primera cita de *Geoica setulosa*
1455 (Passerini, 1860). en la península Ibérica (Hemiptera, Aphididae, Eriosomatidae, Fordini).
1456 *Graellsia* **63**, 143–146.

1457 PÉREZ-LACHAUD, G. & LACHAUD, J.-P. (2014). Arboreal ant colonies as ‘hot-points’ of
1458 cryptic diversity for myrmecophiles: the weaver ant *Camponotus* sp. aff. *textor* and its
1459 interaction network with its associates. *PloS ONE* **9**, e100155.

1460 * PÉREZ-LACHAUD, G., JAHYNY, B.J.B., STÅHLS, G., ROTHERAY, G., DELABIE, J.H.C. &
1461 LACHAUD, J.-P. (2017). Rediscovery and reclassification of the dipteran taxon
1462 *Nothomicrodon* Wheeler, an exclusive endoparasitoid of gyne ant larvae. *Scientific*
1463 *Reports* **7**, 45530.

1464 PÉREZ-LACHAUD, G., ROCHA, F.H., VALLE-MORA, J., HENAUT, Y. & LACHAUD, J.-P. (2019).
1465 Fine-tuned intruder discrimination favors ant parasitoidism. *PloS ONE* **14**, e0210739.

1466 PILOSOV, S., FORTUNA, M. A., VINARSKI, M. V., KORALLO-VINARSKAYA, N. P. & KRASNOV,
1467 B. R. (2013). Temporal dynamics of direct reciprocal and indirect effects in a host–
1468 parasite network. *Journal of Animal Ecology* **82**, 987–996.

1469 PIMM, S.L. (1979). Complexity and stability : another look at MacArthur’s original
1470 hypothesis. *Oikos* **33**, 351–357.

1471 * PLUMARI, M. & JOHARCHI, O. (2017). Rediscovery and redescription of *Laelaps lignicola* G.
1472 & R. Canestrini, a remarkable myrmecophilous mite of the genus *Cosmolaelaps* Berlese
1473 (Acari: Mesostigmata: Laelapidae) from Italy. *Zootaxa* **4232**, 21–40.

- 1474 * POINAR, G. O. (1975). Entomogenous nematodes: a manual and host list of insect-nematode
1475 associations. Brill Archive.
- 1476 POINAR, G. (2012). Nematode parasites and associates of ants: past and present. *Psyche: A*
1477 *Journal Of Entomology* **2012**, 192017.
- 1478 * POLDI, B. (1994). *Strongylognathus pisarskii* species nova (Hymenoptera, Formicidae).
1479 *Memorabilia Zoologica* **48**, 187–191.
- 1480 * PONTIN, A.J. (1963) Further considerations of competition and the ecology of the ants *Lasius*
1481 *Flavus* (F.) and *L. niger* (L.). *Journal of Animal Ecology* **32**, 565–574.
- 1482 POULIN, R. (1995). Phylogeny, ecology, and the richness of parasite communities in vertebrates.
1483 *Ecological Monographs* **65**, 283–302.
- 1484 POULIN, R. & MOUILLOT, D. (2003). Parasite specialization from a phylogenetic perspective:
1485 a new index of host specificity. *Parasitology* **126**, 473–480.
- 1486 POULIN, R. & MOUILLOT, D. (2005). Combining phylogenetic and ecological information into
1487 a new index of host specificity. *Journal of Parasitology* **91**, 511–514.
- 1488 * RADCHENKO, A.G. (1985). Ants of the genus *Strongylognathus* (Hymenoptera: Formicidae)
1489 in the European part of the USSR. *Zoologicheskii zhurnal*. **64**, 1514–1523.
- 1490 * REEMER, M. (2013). Review and phylogenetic evaluation of associations between
1491 Microdontinae (Diptera: Syrphidae) and ants (Hymenoptera: Formicidae). *Psyche: A*
1492 *Journal Of Entomology* **2013**, 538316.
- 1493 * REEMER, M., RENEMA, W., VAN STEENIS, W., ZEEGERS, T., BARENDREGT, A., SMIT, J.T.,
1494 VEEN, M. VAN, VAN STEENIS, J. & VAN DER LEIJ, L. (2009). *De Nederlandse Zweefvliegen*.
1495 Nederlandse Fauna 8. Leiden. Nationaal Natuurhistorisch Museum Naturalis, KNNV
1496 Uitgeverij.

1497 RETTENMEYER, C.W., RETTENMEYER, M.E., JOSEPH, J. & BERGHOFF, S.M. (2010). The largest
 1498 animal association centered on one species: the army ant *Eciton burchellii* and its more
 1499 than 300 associates. *Insectes Sociaux* **58**, 281–292.

1500 RICO-GRAY, V. & OLIVEIRA, P. S. (2007). *The ecology and evolution of ant-plant interactions*.
 1501 University of Chicago Press.

1502 * ROBERTI, D. (1996). Aggiunte alla monografia : Gli Afidi d'Italia. *Entomologica* **30**, 207–
 1503 249.

1504 ROCHA, F.H., LACHAUD, J.-P. & PÉREZ-LACHAUD, G. (2020). Myrmecophilous organisms
 1505 associated with colonies of the ponerine ant *Neoponera villosa* (Hymenoptera:
 1506 Formicidae) nesting in *Aechmea bracteata* bromeliads: a biodiversity
 1507 hotspot. *Myrmecological News* **30**, 73–92.

1508 ROSAS-VALDEZ, R. & DE PEREZ-PONCE DE LEON, G. (2011). Patterns of host specificity
 1509 among the helminth parasite fauna of freshwater siluriforms: Testing the biogeographical
 1510 core parasite fauna hypothesis. *Journal of Parasitology* **97**, 361–363.

1511 * RUANO, F., LENOIR, A., SILVESTRE, M., KHALIL, A. & TINAUT, A. (2018). Chemical profiles
 1512 in *Iberoformica subrufa* and *Formica frontalis*, a new example of temporary host–parasite
 1513 interaction. *Insectes Sociaux* **66**, 223–233.

1514 * RUANO, F. & TINAUT, A. (2005). Mating behaviour in a slave-making ant, *Rossomyrmex*
 1515 *minuchae* (Hymenoptera, Formicidae). *Naturwissenschaften* **92**, 328–331.

1516 * RŮŽIČKA, J. & VÁVRA, J.C. (2009). Interesting records of small carrion beetles (Coleoptera :
 1517 Leiodidae : Cholevinae) from Slovakia. *Klapalekiana* **45**, 233–245.

1518 SAAVEDRA, S., ROHR, R. P., BASCOMPTE, J., GODOY, O., KRAFT, N. J. B., & LEVINE, J. M.
 1519 (2017). A structural approach for understanding multispecies coexistence. *Ecological*
 1520 *Monographs* **87**, 470–486.

- 1521 * SABELLA, G., BEKCHIEV, R. & KOSTOVA, R. (2017). Pselaphinae (Coleoptera Staphylinidae)
 1522 from Sicily and Calabria (Italy). Results from a short entomological expedition in April,
 1523 2015. *Biodiversity Journal* **8**, 185–192.
- 1524 * SABOORI, A., KHAUSTOV, A., HAKIMITABAR, M. & HAJIQANBAR, H. (2009). A new genus and
 1525 species of larval Erythraeinae (Acari: Prostigmata: Erythraeidae) from Ukraine and the
 1526 taxonomic state of Zhangielli. *Zootaxa* **2203**, 22–30.
- 1527 * SANCHEZ-GARCIA, D. & ESPADALER, X. (2017). Una nueva especie parásita social para la
 1528 península Ibérica. *Bothriomyrmex communista santschi*, 1919 (Hymenoptera: Formicidae)
 1529 en España. *Iberomyrmex* **9**, 11–13.
- 1530 SANDERS, I.R. (2003). Preference, specificity and cheating in the arbuscular mycorrhizal
 1531 symbiosis. *Trends in Plant Science* **8**, 143–145.
- 1532 * SANETRA, M. & BUSCHINGER, A. (2000). Phylogenetic relationships among social parasites
 1533 and their hosts in the ant tribe Tetramoriini (Hymenoptera: Formicidae). *European Journal*
 1534 *of Entomology* **97**, 95–117.
- 1535 * SANETRA, M., GÜSTEN, R. & SCHULZ, A. (1999). On the taxonomy and distribution of Italian
 1536 *Tetramorium* species and their social parasites. *Memorie della Societa Entomologica*
 1537 *Italiana* **77**, 317–357.
- 1538 * SANTAMARIA, S. & ESPADALER, X. (2015). *Rickia lenoirii*, a new ectoparasitic species, with
 1539 comments on world Laboulbeniales associated with ants. *Mycoscience* **56**, 224–229.
- 1540 * SCHÄR, S. & NASH, D.R. (2014). Evidence that microgynes of *Myrmica rubra* ants are social
 1541 parasites that attack old host colonies. *Journal of Evolutionary Biology* **27**, 2396–2407.
- 1542 * SCHEMBRI, S. (1984). Notes on the Orthoptera of the Maltese Islands: The genus
 1543 *Myrmecophilus* (Orthoptera: Gryllidae). *The Central Mediterranean Naturalist* **1**, 57–59.

- 1544 * SCHEMBRI, S.P. & COLLINGWOOD, C.A. (1981). A revision of the myrmecofauna of the
 1545 Maltese Islands (Hymenoptera, Formicidae). *Ann. Mus. Civ. Stor. Nat. 'Giacomo Doria'*
 1546 **83**, 417-442.
- 1547 * SCHEURER, S. (1963). Zur Biologie einiger Fichten bewohnender Lachnidenarten. *Zeitschrift*
 1548 *für Angewandte Entomologie* **53**, 153–178.
- 1549 * SCHEURER, S. (1969). Materialien zur Verbreitung der auf *Pinus silvestris* L. lebenden
 1550 Cinarinen im Gebiet der DDR. *Hercynia* **6**, 319–337.
- 1551 * SCHEURER, S. (1991). Ein Beitrag zum Vorkommen und zur Biologie einiger Cinara-Species
 1552 in den Rhodopen (Bulgarien) (Homoptera, Aphidina). *Deutsche Entomologische*
 1553 *Zeitschrift* **38**, 371–378.
- 1554 * SCHLAGHAMERSKÝ, J., KAŠPAR, J., PETRÁKOVÁ, L. & ŠUSTR, V. (2013). Trophobiosis in the
 1555 arboricolous ant *Liometopum microcephalum* (Hymenoptera: Formicidae:
 1556 Dolichoderinae). *European Journal of Entomology* **110**, 231–239.
- 1557 * SCHLICK-STEINER, B.C., STEINER, F.M. & BUSCHINGER, A. (2003). Ameisengrillen
 1558 (*Myrmecophilus* spp.) im wissenschaftlichen Niemandsland Mitteleuropas (Ensifera :
 1559 Myrmecophilidae). *Beiträge zur Entomofaunistik* **4**, 67–73.
- 1560 * SCHLICK-STEINER, B.C., STEINER, F.M., SCHÖDL, S. & SEIFERT, B. (2003). *Lasius austriacus*
 1561 sp. n., a Central European ant related to the invasive species *Lasius neglectus*.
 1562 *Sociobiology* **41**, 725–736.
- 1563 * SCHMID, U. (2004). *Microdon rhenanus* and *Microdon eggeri* var. major (Diptera, Syrphidae)
 1564 revisited. *Volucella* **7**, 111–124.
- 1565 * SCHMIDT, C. & SCHMITT, T. (2012). Vorkommen der Ameisengrille (*Myrmecophilus*
 1566 *acervorum* [PANZER, 1799]) im Bochumer Raum. *Jahresberichte Bochumer*

- 1567 *Botanischen Vereins* **3**, 25–32.
- 1568 * SCHMIDT, H. (1952). Zur Kenntnis der Wurzellaus *Paracletus cimiciformis* v. d. Heyd., ein
1569 parasitär angepaßter Ameisengast. *Zeitschrift für Parasitenkunde* **15**, 283–289.
- 1570 SCHOENER, T.W. (1989). Food webs from the small to the large: the Robert H. MacArthur
1571 award lecture. *Ecology* **70**, 1559–1589.
- 1572 SCHÖNROGGE, K., BARR, B., WARDLAW, J.C., NAPPER, E., GARDNER, M.G., BREEN, J.,
1573 ELMES, G.W. & THOMAS, J.A. (2002). When rare species become endangered: cryptic
1574 speciation in myrmecophilous hoverflies. *Biological Journal of the Linnean Society* **75**,
1575 291–300.
- 1576 * SEIFERT, B. (2000). A taxonomic revision of the ant subgenus *Coptoformica* Mueller, 1923
1577 (Hymenoptera, Formicidae). *Zoosystema* **22**, 517–568.
- 1578 SEIFERT, B. (2007). *Die Ameisen Mittel- und Nordeuropas*. Iuta Verlags- und
1579 Vertriebsgesellschaft, Görlitz.
- 1580 * SKORUPSKI, M. & GWIAZDOWICZ, D.J.G. (2002). Mites (Acari, Gamasida) inhabiting ant nests
1581 in the Pieniny Mountains. *Pieniny – Przyroda i Człowiek* **7**, 105–107.
- 1582 * SLIPINSKI, S.A. & BURAKOWSKI, B. (1988). A review of the genus *Rhopalocerus* W.
1583 Redtenbacher of the world. *Annales Zoologici* **42**, 75–118.
- 1584 * SLOGGETT, J.J. & MAJERUS, M.E.N. (2000). Aphid-mediated coexistence of ladybirds
1585 (Coleoptera: Coccinellidae) and the wood ant *Formica rufa*: seasonal effects, interspecific
1586 variability and the evolution of a coccinellid myrmecophile. *Oikos* **89**, 345–359.
- 1587 * SOLDATI, F. & SOLDATI, L. (2000). A propos d' *Oochrotus unicolor* Lucas, 1852 (Coleoptera
1588 Tenebrionidae). *Bulletin de la Société Linnéenne de Bordeaux* **28**, 181–187.
- 1589 SOLÉ, R. V & BASCOMPTE, J. (2006). *Self-Organization of Complex Ecosystems (MPB-42)*.

- 1590 Princeton University Press.
- 1591 SOLÉ, R. V. & MONTOYA, J.M. (2001). Complexity and fragility in ecological networks.
- 1592 *Proceedings of the Royal Society B: Biological Sciences* **268**, 2039–2045.
- 1593 * SOLODOVNIKOV, I.A. & PLISKEVICH, E.S. (2014). Comparison of species composition of the
- 1594 community myrmecophilous coleopteran (Insecta, Coleoptera) in the nest *Formica*
- 1595 *exsecta*, *Formica rufa*, *Formica polyctena* in the Belarus Lakeland. *The scientific-*
- 1596 *theoretical and information-methodical journal of the Belarusian Republican Foundation*
- 1597 *for Fundamental Research* **4**, 37–46.
- 1598 SPAAK, J. & DE LAENDER, F. (2020). Intuitive and broadly applicable definitions of niche and
- 1599 fitness differences. *Ecology Letters*, doi: <https://doi.org/10.1111/ele.13511>.
- 1600 * STADLER, B. & DIXON, A.F.G. (2008). *Mutualism: Ants and Their Insect Partners*. Cambridge
- 1601 University Press, Cambridge.
- 1602 * STADLER, B., KINDLMANN, P., SMILAUER, P. & FIEDLER, K. (2003). A comparative analysis
- 1603 of morphological and ecological characters of European aphids and lycaenids in relation
- 1604 to ant attendance. *Oecologia* **135**, 422–430.
- 1605 * STALLING, T. (2013a) A new species of ant-loving cricket from Mallorca, Balearic Islands,
- 1606 Spain (Orthoptera, Myrmecophilidae). *Graellsia* **69**, 153–156.
- 1607 * STALLING, T. (2013b) First records of the ant-loving cricket *Myrmecophilus acervorum*
- 1608 (Panzer, [1799]) from Armenia and comments on *Myrmecophilus hirticaudus* Fischer von
- 1609 Waldheim, 1846 (Orthoptera: Myrmecophilidae). *Biological Journal of Armenia* **2**, 120–
- 1610 122.
- 1611 * STALLING, T. (2013c) *Myrmecophilus balcanicus*, a new species of ant-loving cricket from
- 1612 the Former Yugoslav Republic of Macedonia, with notes on the synonymy of
- 1613 *Myrmecophilus zorae*. *Entomologia Hellenica* **22**, 29–34.

- 1614 * STALLING, T. (2015). New records of the ant-loving cricket *Myrmecophilus fuscus* Stalling,
1615 2013 from the Mediterranean Basin (Orthoptera: Myrmecophilidae). *Articulata* **30**, 105–
1616 108.
- 1617 * STALLING, T. (2017a). A new species of ant-loving cricket *Myrmecophilus* Berthold, 1827
1618 from Cyprus (Orthoptera: Myrmecophilidae). *Zoology in the Middle East* **63**, 63–67.
- 1619 * STALLING, T. (2017b). *Myrmecophilus gallicus*, a new species of ant cricket from France
1620 (Orthoptera: Myrmecophilidae). *Articulata* **32**, 1-7.
- 1621 * STALLING, T. & BIRRER, S. (2013). Identification of the ant-loving crickets, *Myrmecophilus*
1622 Berthold, 1827 (Orthoptera: Myrmecophilidae), in Central Europe and the northern
1623 Mediterranean Basin. *Articulata* **28**, 1–11.
- 1624 * STANIEC, B. & ZAGAJA, M. (2008). Rove-beetles (Coleoptera, Staphylinidae) of ant nests of
1625 the vicinities of Leżajsk. *Annales Universitatis Mariae Curie-Sklodowska Lublin - Polonia*
1626 **63**, 111–127.
- 1627 * STAVERLØKK, A. & ØDEGAARD, F. (2016). New records of parasitic Hymenoptera associated
1628 with ants, including four new species to Norway. *Norwegian Journal of Entomology* **63**,
1629 188–196.
- 1630 STEPHENS, P.R., ALTIZER, S., SMITH, K.F., AGUIRRE, A.A., BROWN, J.H., BUDISCHAK, S.A.,
1631 BYERS, J.E., DALLAS, T.A., DAVIES, T.J., DRAKE, J.M., EZENWA, V.O., FARRELL, M.J.,
1632 GITTLEMAN, J.L., HAN, B.A., HUANG, S., ET AL. (2016). The macroecology of infectious
1633 diseases: a new perspective on global-scale drivers of pathogen distributions and impacts.
1634 *Ecology Letters* **19**, 1159–1171.
- 1635 STOEFFLER, M., TOLASCH, T. & STEIDLE, J.L.M. (2011). Three beetles—three concepts.
1636 Different defensive strategies of congeneric myrmecophilous beetles. *Behavioral*

- 1637 *Ecology and Sociobiology* **65**, 1605–1613.
- 1638 * STUART, R. J., & ALLOWAY, T. M. (1988). Aberrant yellow ants: North American
 1639 *Leptothorax* species as intermediate hosts of cestodes, pp. 537-545 in J. Trager [ed.],
 1640 Advances in myrmecology. E. J. Brill, Leiden.
- 1641 * SUAY-CANO, V.A., TINAUT, A. & SELFA, J. (2002). Las hormigas (Hymenoptera, Formicidae)
 1642 asociadas a pulgones (Hemiptera, Aphididae) en la provincia de Valencia. *Graellsia* **58**,
 1643 21–37.
- 1644 * SUDD, J.H. & SUDD, M.E. (1985). Seasonal changes in the response of wood ants (*Formica*
 1645 *lugubris*) to sucrose baits. *Ecological Entomology* **10**, 89–97.
- 1646 * SZATHMÁRY, E., HALTRICH, A. & TARTALLY, A. (2005). Data to the knowledge of the Lachnid
 1647 fauna (Homoptera: Lachnidae) of Hungary. *Acta Phytopathologica et Entomologica*
 1648 *Hungarica* **40**, 403–408.
- 1649 * SZELEGIEWICZ, H. (1959). Redescription of *Lachnus* (*Shizodryobius*) *longirostris* (Mordv.,
 1650 1901), with notes on the type specimens. *Annales Zoologici* **13**, 191–220.
- 1651 * SZELEGIEWICZ, H. (1961). Die polnischen Arten der gattung *Chaitophorus* Koch s. lat.
 1652 (Homoptera, Aphididae). *Annales Zoologici* **7**, 229–352.
- 1653 TARTALLY, A., THOMAS, J.A., ANTON, C., BALLETO, E., BARBERO, F., BONELLI, S., BRÄU,
 1654 M., CASACCI, L. P., CSÖSZ, S., CZEKES, Z., DOLEK, M., DZIEKAŃSKA, I., ELMES, G.,
 1655 FÜRST, M.A., GLINKA, U., ET AL. (2019). Patterns of host use by brood parasitic
 1656 *Maculinea* butterflies across Europe. *Philosophical Transactions of the Royal Society B*
 1657 **374**, 20180202.
- 1658 * TELNOV, D. (2008). Beetles (Coleoptera) living in the nests of *Lasius fuliginosus* (Latreille,
 1659 1798). (Hymenoptera : Formicidae) in Latvia. *Latvijas entomologs* **46**, 70–71.

- 1660 THOMAS, J.A., SCHÖNROGGE, K. & ELMES, G.W. (2005). Specializations and host associations
1661 of social parasites of ants. In *Insect Evolutionary Ecology* (eds M.D.E. FELLOWES, G.J.
1662 HOLLOWAY & J. ROLFF), pp. 479–518. Royal Entomological Society, CABI Publishing,
1663 UK.
- 1664 THRALL, P.H., HOCHBERG, M.E., BURDON, J.J. & BEVER, J.D. (2007). Coevolution of
1665 symbiotic mutualists and parasites in a community context. *Trends in Ecology and*
1666 *Evolution* **22**, 120–126.
- 1667
- 1668 * TICHÁ, K. (2006). The first record of *Gursaultia giguntonympha* (Acari : Acaridae) an
1669 *Temnothorax crassispinus* (Hymenoptera : Formicidae). *Acta rerum naturalium* **2**, 49–50.
- 1670 TILMAN, D. (1982). *Resource Competition and Community Structure (MBP-17)*. Princeton
1671 University Press.
- 1672 * TIZADO, E.J., TINAUT, A. & NIETO NARIA, J.M. (1993). Relationships between ants and aphids
1673 in the province of Leon (Spain) (Hym: Formicidae; Hom : Aphididae). *Vie Milieu* **43**, 63–
1674 68.
- 1675 * TORKA, V. (1905). *Tettigometra obliqua* Panz. *Zeitschrift für wissenschaftliche*
1676 *Insektenbiologie* **1**, 451–455.
- 1677 * TRACH, V. & KHAUSTOV, A. (2011). A Myrmecophilous Mite *Myrmozercon tauricus* sp. n.
1678 of the family Laelapidae (Acari, Mesostigmata) from Ukraine. *Vestnik Zoologii* **45**, 23–
1679 27.
- 1680 TRØJELSGAARD, K. & OLESEN, J.M. (2013). Macroecology of pollination networks. *Global*
1681 *Ecology and Biogeography* **22**, 149–162.
- 1682 TYKARSKI, P. (2017). Coleoptera Poloniae - Information System about Beetles of Poland.
1683 <http://coleoptera.ksib.pl/?l=en>.
- 1684 * UPPSTROM, K.A. (2010). *Mites (Acari) associated with the ants (Formicidae) of Ohio and the*

1685 *harvester* Ant, *Messor pergandei*, of Arizona. Master thesis, Ohio State University.

1686 * URBANO GRANERO, J.M. (2003). Primera cita para Huelva (Andalucía) de *Amorphocephala*

1687 *coronata* (Germar, 1817). (Coleoptera : Brenthidae). *Boletín de la SAE* **7**, 16–18.

1688 * VAN ACHTERBERG, C. & ORTIZ DE ZUGASTI CARRÓN, N.F. (2016). Revision of the genus

1689 *Paralipsis* Foerster, 1863 (Hymenoptera, Braconidae), with the description of two new

1690 species. *ZooKeys* **2016**, 25–39.

1691 VALDOVINOS, F.S. (2019). Mutualistic networks: moving closer to a predictive

1692 theory. *Ecology letters* **22**, 1517–1534.

1693 * VANTAUX, A., ROUX, O., MAGRO, A. & ORIVEL, J. (2012). Evolutionary perspectives on

1694 myrmecophily in ladybirds. *Psyche: A Journal Of Entomology* **2012**, 591570.

1695 * VÁVRA, J.C. (2015). Review of the subgenus *Attumbrinus* (Coleoptera : Leiodidae :

1696 Cholevinae : *Philomessor*), with description of *Philomessor lackneri* sp . nov . from

1697 Morocco. *Acta Entomologica Musei Nationalis Pragae* **55**, 85–103.

1698 VAZQUEZ, D.P., CHACOFF, N.P. & CAGNOLO, L. (2009). Evaluating multiple determinants of

1699 the structure of plant-animal mutualistic networks. *Ecology* **90**, 2039–2046.

1700 VELLEND, M. (2016). *The Theory of Ecological Communities (MPB-57)*. Princeton University

1701 Press.

1702 * VEPSÄLÄINEN, K. & CZECHOWSKI, W. (2014). Against the odds of the ant competition

1703 hierarchy: Submissive *Myrmica rugulosa* block access of the dominant *Lasius fuliginosus*

1704 to its aphids. *Insectes Sociaux* **61**, 89–93.

1705 VERESOGLOU, S.D. & RILLIG, M.C. (2014). Do closely related plants host similar arbuscular

1706 mycorrhizal fungal communities ? A meta-analysis. *Plant and Soil* **377**, 395–406.

1707 VILLA, S.M., LE BOHEC, C., KOOP, J.A.H., PROCTOR, H.C. & CLAYTON, D.H. (2013).

1708 Diversity of feather mites (Acari: Astigmata) on Darwin’s finches. *Journal of*

- 1709 *Parasitology* **99**, 756–762.
- 1710 * VIÑOLAS, A. (1981). Tenebrionidos de la comarca del Segria y de las zonas halofilas de Los
1711 Monegros (Coleoptera). *II Sessió conjunta d'entomologia*, 67–72.
- 1712 * VIÑOLAS, A., ESPADALER, X., GARCÍA, F. & ROIG, X. (2008). *Lomechusa emarginata*
1713 (Paykull, 1789). (Coleoptera: Staphylinidae: Aleocharinae), nueva para Catalunya, en nido
1714 de *Myrmica spinosior* Santschi, 1931 (Hymenoptera: Formicidae). *Heteropterus Rev*
1715 *Entomol* **8**, 207–210.
- 1716 * VITZTHUM, H. (1917). Neue myrmecophile Milben. *Zoologischer Anzeiger* **50**, 188–192.
- 1717 * VÖLKL, W. (1995). Behavioral and morphological adaptations of the coccinellid, *Platynaspis*
1718 *luteorubra* for exploiting ant-attended resources (Coleoptera: Coccinellidae). *Journal of*
1719 *Insect Behavior* **8**, 653–670.
- 1720 VON BEEREN, C., MARUYAMA, M. & KRONAUER, D.J.C. (2015). Cryptic diversity, high host
1721 specificity and reproductive synchronization in army ant-associated *Vatesus* beetles.
1722 *Molecular Ecology* **25**, 990–1005.
- 1723 * WAGNER, H.C., KOSCHUH, A., SCHATZ, I. & STALLING, T. (2012). Die Myrmekophilen einer
1724 Lawinenrinne im Nationalpark Gesäuse (Steiermark). *Abhandlungen der Zoologisch-*
1725 *Botanischen Gesellschaft in Österreich*. **38**, 147–161.
- 1726 WAGNER, K., MENDIETA-LEIVA, G. & ZOTZ, G. (2015). Host specificity in vascular epiphytes:
1727 A review of methodology, empirical evidence and potential mechanisms. *AoB PLANTS*
1728 **7**, 1–25.
- 1729 * WAHAB, A. (1962). Untersuchungen über Nematoden in den Drüsen des Kopfes der Ameisen
1730 (Formicidae). *Zeitschrift für Morphologie und Ökologie der Tiere* **52**, 33–92.
- 1731 WASMANN, E. (1894). *Kritisches Verzeichniss der myrmekophilen und termitophilen*
1732 *Arthropoden Mit Angabe der Lebensweise und mit Beschreibung neuer Arten*. Berlin: F.

- 1733 L. Dames, xv.
- 1734 * WASMANN, E. (1899a) Weitere nachträge zum verzeichniss der Ameisengäste von
1735 Holländisch Limburg. *Tijdschrift Voor Entomologie* **42**, 158–171.
- 1736 * WASMANN, E. (1899b) Zur Kenntniss der bosnischen Myrmekophilen und Ameisen.
1737 *Wissenschaftliche Mitteilungen aus Bosnien Hercegovina* **6**, 767–772.
- 1738 * WEBER, S. (2013). Waldameisen-Lachniden-Trophobie und Qualität des Honigtaus von
1739 Fichtenrindenläusen (Sternorrhyncha, Lachnidae). Thesis. Technische Universität
1740 München.
- 1741 * WEISSFLOG, A., MASCHWITZ, U., SEEBAUER, S., DISNEY, R.H.L., SEIFERT, B. & WITTE, V.
1742 (2008). Studies on European ant decapitating flies (Diptera: Phoridae): II. Observations
1743 that contradict the reported catholicity of host choice by *Pseudacteon formicarum*.
1744 *Sociobiology* **51**, 87–94.
- 1745 WERNER, E. E., & GILLIAM, J. F. (1984). The ontogenetic niche and species interactions in
1746 size-structured populations. *Annual Review of Ecology and Systematics* **15**, 393–425.
- 1747 * WITEK, M., BARBERO, F. & MARKÓ, B. (2014). *Myrmica* ants host highly diverse parasitic
1748 communities: from social parasites to microbes. *Insectes Sociaux* **61**, 307–323.
- 1749 * WITEK, M., PATRICELLI, D., CASACCI, L. PIETRO, BARBERO, F., BALLETO, E. & BONELLI, S.
1750 (2011). Notes on the biology and host ant specificity of the myrmecophilous syrphid fly
1751 *Microdon major* (diptera: Syrphidae), a social parasite of *Formica* ants (Hymenoptera:
1752 Formicidae). *Sociobiology* **57**, 261–269.
- 1753 * WITHERS, P. (1994). Quelques Scatopsidae de la région lyonnaise, dont trois espèces
1754 nouvelles pour la France (Diptera Scatopsidae). *Bulletin mensuel de la Société Linnéenne
1755 de Lyon* **63**, 335–336.

- 1756 * WURST, E. (2001). The Life Cycle of *Lemanniella minotauri* n. sp. and the erection of the
1757 new family Lemanniellidae (Acari: Astigmata). *Stuttgarter Beiträge zur Naturkunde* **621**,
1758 1–34.
- 1759 * YÉLAMOS, T. (1992). Revision del genero *Eretmotus* Lacordaire, 1854 (Coleoptera,
1760 Histeridae). *Eos* **68**, 8–27.
- 1761 * YÉLAMOS, T. (1995). Revision of the genus *Sternocoelis* Lewis, 1888 (Coleoptera:
1762 Histeridae), with a proposed phylogeny. *Revue Suisse de Zoologie* **102**, 113–174.
- 1763 * YÉLAMOS, T., BOLAÑO, S. & BLANCO VILLERO, J.M. (2010). Nuevas aportaciones al catálogo
1764 de maláquidos (Coleoptera, Malachiidae) de la Sierra de Tudía (Badajoz, Extremadura,
1765 España). *Boletín de la Sociedad Entomológica Aragonesa (S.E.A.)* **47**, 418.
- 1766 ZAGAJA, M. & STANIEC, B. (2015). *Thiasophila szujekii* sp. n. (Coleoptera, Staphylinidae,
1767 Aleocharinae)-a cryptic species associated with *Formica truncorum* in Poland. *Zootaxa*
1768 **3955**, 417–426.
- 1769 * ZWÖLFER, H. (1958). Zur Systematik, Biologie und Ökologie unterirdisch lebender Aphiden
1770 (Homoptera, Aphidoidea) (Anoeciinae, Tetraneurini, Pemphigini und Fordinae). Teil II
1771 Tetraneurini und Pemphigini. *Zeitschrift für angewandte Entomologie*, 528–575.

1772

1773 **IX. SUPPORTING INFORMATION**

1774 **Appendix S1.** Host–symbiont matrix listing the associations between ants and ant symbionts
1775 in Europe. Ant species are listed in columns and symbionts in rows. 1, interaction recorded; 0,
1776 no interaction recorded.

1777 **Appendix S2.** List of literature used to reconstruct the host–symbiont matrix in Appendix S1.

1778 **Appendix S3.** Host–symbiont matrix for ant hosts with known phylogeny.

1779 **Appendix S4.** Predictor values of host ant species.

1780 **Fig. S1.** Correlation between the number of described European species in an ant genus and
1781 the number of associated symbionts.

1782 **Fig. S2.** Number of host species per symbiont type.

1783 **Fig. S3.** Mean \pm SE taxonomic distance of targeted host species for different types of ant
1784 symbionts. Symbionts with a single host species were omitted from this analysis. Letter codes
1785 refer to Tukey *post-hoc* test. Symbiont types with no letters in common are significantly
1786 different ($P < 0.05$).

1787 **Fig. S4.** Mean \pm SE taxonomic distance of targeted host species for different types of ant
1788 symbionts. Taxonomic distance of symbionts with a single host species was set at 1.

1789

1790

FIGURE LEGENDS

Fig. 1. Distribution of different types of ant symbionts. Total number of symbionts $N = 722$, number of symbionts per type given in parentheses. Trophobionts are mutualistic, social parasites and helminths are strictly parasitic. Fungi are parasitic, except for the mutualistic species *Cladosporium myrmecophilum*. Myrmecophiles range from commensals to parasites and include parasitoid wasps and flies. Note that three mutualistic Lepidoptera species are classified as trophobionts and six parasitic and commensalistic Lepidoptera species as myrmecophiles.

Fig. 2. Ant–symbiont network displaying the proportional distribution of symbionts across the European ant genera. A multilevel pie chart is given for each ant genus. The size of the outer pie chart corresponds to the total number of associated symbionts (circle size legend in right corner). The size of the inner pie chart is related to the number of symbionts that are not shared with other ant genera (unique symbionts). The proportional distribution of the five types of ant symbionts (see Fig. 1) is given for all associated symbionts (colour segments in outer pie charts) and for the symbionts that are not shared with other genera (colour segments in inner pie charts). The relative proportion of unique symbionts can be deduced by the relative size of the inner circle to the outer circle. The genera are organized in four groups, corresponding to the ant subfamily to which they belong. The genera are connected with lines, of which the width is directly proportional to the number of shared symbionts. The font size of a genus is proportional to its number of described species in Europe.

Fig. 3. Mean \pm SE number of host species per symbiont type, controlled for sampling effort. Letter codes refer to Tukey *post-hoc* test. Species with no letters in common are significantly different ($P < 0.05$).

1816

1817 **Fig. 4.** Mean \pm SE phylogenetic distance of targeted host species (based on the phylogenetic
1818 tree of Arnan *et al.*, 2017) for different types of ant symbionts. Letter codes refer to *post-hoc*
1819 test. Symbiont types with no letters in common are significantly different ($P < 0.05$).

1820


1821 **Fig. 5.** Ranking of the predictors from the five PGLS models by the corrected Akaike
1822 information criterion (AICc). The change in AICc (Δ AICc) when adding or removing a
1823 variable from the most optimal model is compared. Predictors included in the most optimal
1824 model are removed (Δ AICc positive), whereas those not included are added (Δ AICc negative)
1825 to the best model (lowest AICc). The ranking is given for the five PGLS analyses, i.e. with
1826 dependent variable the number of symbionts (best model: \sim sample effort + colony size +
1827 distribution + habitat), myrmecophiles (best model: \sim sample effort + colony size +
1828 distribution + habitat + worker size), trophobionts (best model: \sim sample effort + habitat),
1829 social parasites (best model: \sim distribution range) and helminths (best model: \sim sample effort
1830 + habitat), respectively. Note that myrmecophiles, trophobionts, social parasites and
1831 helminths are four subsets of all ant-associated symbionts.

1832


1833 **Fig. 6.** Relative importance of the significant predictor matrices explaining the dissimilarity in
1834 symbiont communities across different ant species. Rankings are given for predictors
1835 explaining overall dissimilarity (1–similarity) in symbiont composition, and for dissimilarity
1836 in subsets of symbiont composition: myrmecophiles, trophobionts, social parasites, helminths
1837 and parasitic fungi, respectively. Note that myrmecophiles, trophobionts, social parasites,
1838 helminths and parasitic fungi are subsets of all ant-associated symbionts. The allocated
1839 contribution (sequential R^2) of the different distance matrices (indicated with Δ) or the matrix
1840 capturing the combined sample effort of a pair of host species to the explained variation of the

1841 MRM models is estimated with the lmg metric. The error bars are 95% confidence intervals
1842 produced using 1000 bootstrap replicates. The combined sample effort of a pair of host
1843 species was negatively correlated with their dissimilarity in symbiont composition. The
1844 dissimilarity matrices of all other predictors were positively correlated with dissimilarity in
1845 symbiont composition. Significance levels of the predictors were tested with a permutation
1846 test ($N = 9999$; ***, $P < 0.001$; **, $P < 0.01$; *, $P < 0.05$; ·, $P < 0.10$).

1847



1850
1851 Fig. 1



1852

1853 Fig.2




Fig. 3






Fig. 4



1860

1861 Fig. 5

1862



1863
1864
1865
1866

Fig. 6

1868 Table 1. Overview of the different types of ant symbionts found in Europe.

Symbiont type	General description	Representatives and strategies	Ref.
MYRMECOPHILE	A diverse group of arthropods that mostly live inside ant nests. The life strategies of these organisms range from commensalism to specialized parasitism; there are no mutualists in this group. Representatives in many arthropod orders, but beetles and mites are the most diverse. Also known as ant guests.	<ul style="list-style-type: none"> - <u>unspecialized myrmecophiles</u>: poorly integrated in the colony and may provoke aggression. Very similar in behaviour and morphology to free-living relatives. Do not rely on advanced chemical deception. - <u>specialized myrmecophiles</u>: accepted in the colony by being groomed, fed or transported. Typically possess advanced glandular structures (trichomes) or specialized behaviour to deceive the host. - <u>myrmecophilous parasitoids</u>: wasps and flies of which the larvae parasitize ant workers, ant larvae or possibly other myrmecophiles. Eventually kill the host. Adult parasitoids do not live in the nest. 	1–6
TROPHOBIONT	Mutualistic arthropods that provide sugary honeydew in exchange for protection and hygienic services. Mostly live outside the nest.	<ul style="list-style-type: none"> - <u>Hemiptera</u>: aphids, scale insects and planthoppers. - <u>Lepidoptera</u>: mutualistic caterpillars permanently living outside the nest. 	1
SOCIAL PARASITE	A group of ants that parasitize other ant species.	<ul style="list-style-type: none"> - <u>xenobiosis</u>: parasitic ants that construct a nest inside other ant nests, but raise their own brood. - <u>temporary parasitism</u>: a parasitic queen usurps a host colony and exploits the host work force to establish her own colony. Parasite workers gradually substitute the host worker force. - <u>dulosis (slavery)</u>: a parasite colony is established as in temporary parasitism, but here the workers of the parasitic species will raid pupae of other ant species. Workers which will emerge from these pupae will do most of the tasks in the colony. - <u>inquilinism</u>: parasitic queens permanently exploit a host colony. The parasitic queen produces only sexuals, not workers. 	7
HELMINTH	Endoparasitic worms	<ul style="list-style-type: none"> - <u>nematodes (Nematoda)</u>: the juveniles (dauers) of some groups, such as the Rhabditidae and Diplogastridae, live in the postpharyngeal glands of their ant host and are considered weak parasites. Mermithid nematodes develop in the haemocoel of the ant, may cause morphological changes in the host, and ultimately kill the host upon emergence. Other hosts, such as oligochaetes, may be necessary to complete the life cycle of mermithids. 	8–9

		<p>- <u>flukes (Platyhelminthes: Trematoda)</u>: <i>Dicrocoelium</i> is a trematode whose definitive hosts are grassland vertebrates. The eggs are released along with the faeces and eaten by snails. Ants serve as the second intermediate host of the parasite and become infected by ingesting snail slime. The parasite induces behavioural changes in the ant which then climb to the top of grass stems where they are ingested by the definitive host.</p> <p>- <u>tapeworms (Platyhelminthes: Cestoda)</u>: cestodes are infamous parasites that live in the digestive tract of vertebrates. These tapeworms have a life cycle with multiple hosts and ants may serve as an intermediate host.</p>	
FUNGUS	A diverse group of mainly ant-specific ecto- and endoparasites.	<p>- <u>parasitic fungi</u>: Laboulbeniales are ectoparasites that do not kill their host. They produce a multicellular thallus externally attached to the integument of the host ant. <i>Myrmicinosporidium durum</i> is an endoparasitic fungus which ultimately kills its host. <i>Pandora formicae</i> is a well-known entomopathogenic fungus, that manipulates its ant host to climb the vegetation. The ant attaches itself to the distal part of leaves with its mandibles and dies of the infection.</p> <p>- <u>mutualistic fungi</u>: <i>Cladosporium myrmecophilum</i> provides stability and structure to the carton nests of <i>Lasius fuliginosus</i>.</p>	10, 11

1869 1, Hölldobler & Wilson (1990); 2, Kronauer & Pierce (2011); 3, Elizalde *et al.* (2018); 4, Pérez-Lachaud *et al.* (2019); 5, Parmentier (2020); 6, Kistner (1982); 7, Buschinger
1870 (2009); 8, Poinar (2012); 9, Demartin (2018); 10, Espadaler & Santamaria (2012); 11, Maschwitz & Hölldobler (1970).

1871 Table 2. Estimates for the predictors of the top-ranked PGLS analyses ($\Delta\text{AICc} < 2$). The subset of best models is given for the analyses with total
1872 number of (1) symbionts, (2) myrmecophiles, (3) trophobionts (4) social parasites and (5) helminths as dependent variable. Significant estimates
1873 indicated in bold (***, $P < 0.001$; **, $P < 0.01$; *, $P < 0.05$; , $P < 0.10$).
1874

	intercept	colony size	sample effort	biogeo graphy	distribution	habitat	sympatric ants	nest type	worker size	d.f.	AICc	weight	lambda
all symbionts													
	6.48	0.82***	0.89**		0.70***	+				7	375.3	0.33	0.54
	7.93	0.71***	0.84**		0.74***	+		+		10	375.5	0.30	0.52
	6.85	0.82***	0.82**		0.77***	+			0.31	8	376.0	0.22	0.44
	8.20	0.70***	0.77**		0.80***	+		+	0.26	11	376.8	0.15	0.41
myrmecophiles													
	6.29	0.81***	0.43		0.95***	+			0.45*	8	299.0	0.31	0.00
	6.64	0.94***			1.17***	+			0.46*	7	299.1	0.29	0.00
	7.95	0.75***			1.20***	+		+	0.44*	10	300.4	0.15	0.00
	7.56	0.57***	0.66*	+	0.80***	+			0.57*	12	300.7	0.13	0.32
	6.21	0.81***	0.45	+	1.10***	+	-0.22		0.42*	9	300.9	0.12	0.00
trophobionts													
	2.93		0.77***			+				5	131.2	0.42	0.73
	3.14		0.64***		0.20***	+				6	132.3	0.24	0.60
	3.52	0.22**	0.42*		0.31***	+				7	133.0	0.18	0.36
	3.06	0.11**	0.26***			+				6	133.2	0.16	0.68
social parasites													
	1.34				0.27**					2	75.5	0.23	1.00
	1.19		0.26**							2	75.9	0.19	1.00
	1.25						0.22**			2	76.4	0.15	0.83
	1.30				0.17**		0.11			3	76.7	0.12	0.94
	1.27		0.13		0.16**					3	76.9	0.11	1.00
	1.21		0.16**				0.11			3	76.9	0.11	0.93
	1.29				0.26**				-0.10	3	77.3	0.09	0.98
helminths													
	1.81		0.14*			+				5	31.7	0.18	0.91
	1.83		0.11*			+	0.10			6	31.8	0.16	0.88
	2.01					+	0.13*			5	31.9	0.16	0.85
	1.87		0.11		0.11**	+				6	32.0	0.15	0.88
	2.07				0.14**	+				5	32.1	0.14	0.87

1875

2.07			***		4	33.5	0.07	0.88
1.82	-0.06	0.16*	***		6	33.5	0.07	0.94
1.83	-0.07	0.14*	***	0.10	7	33.5	0.07	0.94
